

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/505, C07D 239/86, 471/04		A1	(11) International Publication Number: WO 94/21259 (43) International Publication Date: 29 September 1994 (29.09.94)
(21) International Application Number: PCT/US94/02834 (22) International Filing Date: 16 March 1994 (16.03.94) (30) Priority Data: 033,595 19 March 1993 (19.03.93) US (60) Parent Application or Grant (63) Related by Continuation US 033,595 (CON) Filed on 19 March 1993 (19.03.93) (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BAGLEY, Scott, W. [US/US]; Apartment 38, 128 North Street, Groton, CT 06340 (US). CHAKRAVARTY, Prasun, K. [IN/US]; 16 Churchill Road, Edison, NJ 08820 (US). CHEN, Anna [US/US]; 852 Madison Avenue, Rahway, NJ 07065 (US). DHANOA, Daljit, S. [CA/US]; 258 Sunset Key, Secaucus, NJ 07094 (US). FITCH, Kenneth, J. [US/US]; 105 Besler Avenue, Cranford, NJ 07016 (US). GREENLEE, William,		J. [US/US]; 115 Hemrick Avenue, Teaneck, NJ 07666 (US). NAYLOR, Elizabeth, M. [GB/US]; 2219 North Avenue, #7, Scotch Plains, NJ 07076 (US). TATA, James, R. [US/US]; 25 Faulkner Drive, Westfield, NJ 07090 (US). WALSH, Thomas, F. [US/US]; 127 Marion Avenue, Westfield, NJ 07090 (US). WILLIAMS, David, L., Jr. [US/US]; 37 Sugarhill Drive, Telford, PA 18969 (US). (74) Agent: CAMARA, Valerie, J.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TT, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
		Published With international search report.	

(54) Title: QUINAZOLINONES SUBSTITUTED WITH PHENOXYPHENYLACETIC ACID DERIVATIVES



(57) Abstract

Phenoxyphenylacetic acids and derivatives of general structural formula (I) have endothelin antagonist activity and are therefore useful in treating cardiovascular disorders, such as hypertension, postischemic renal failure, vasospasm, cerebral and cardiac ischemia, myocardial infarction, inflammatory diseases, Raynaud's disease and endotoxic shock, and asthma.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

TITLE OF THE INVENTION
QUINAZOLINONES SUBSTITUTED WITH
PHENOXYPHENYLACETIC ACID DERIVATIVES

5 SUMMARY OF THE INVENTION

This invention is concerned with non-peptidic endothelin receptor antagonists and their method of use. The compounds of the present invention are therapeutic agents particularly useful for the treatment of asthma, hypertension, pulmonary hypertension, arteriosclerosis, congestive heart failure, renal failure, particularly post-ischemic renal failure, cyclosporin nephrotoxicity, vasospasm, vascular restenosis, cerebral and cardiac ischemia and other ischemic states, myocardial infarction, Raynaud's disease, inflammatory bowel diseases, including Crohn's disease and ulcerative colitis, as well as other inflammatory diseases, or endotoxic shock caused by or associated with endothelin.

BACKGROUND OF THE INVENTION

Endothelin is a 21-amino acid peptide produced by endothelial cells. The peptide is secreted not only by endothelial cells but also by tracheal epithelial cells or from kidney cells. Endothelin (ET-1) has a potent vasoconstrictor effect. The vasoconstricting effect is caused by the binding of endothelin to its receptor on the vascular smooth muscle cells.¹⁻³

Endothelin-1 (ET-1) is one of three recently identified potent vasoconstricting peptides which also includes endothelin-2 (ET-2) and endothelin-3 (ET-3) which differ from ET-1 by two and six amino acids, respectively.⁴

Increased levels of endothelin are found in the blood of patients with essential hypertension, acute myocardial infarction, pulmonary hypertension, Raynaud's disease or atherosclerosis or in the washing fluids of the respiratory tract of patients with asthma compared to normal levels.⁵⁻⁸

- 2 -

An experimental model of cerebral vasospasm and a second model of acute renal failure have led to the conclusion that endothelin is one of the mediators causing cerebral vasospasm following a subarachnoid hemorrhage, and renal failure.⁹⁻¹⁰

5 Endothelin was also found to control the release of many physiological substances such as renin, atrial natriuretic peptide, endothelium-derived relaxing factor (EDRF), thromboxane A₂,¹⁴, prostacyclin, norepinephrine, angiotensin II and substance P.¹¹⁻¹⁶ Further, endothelin causes contraction of the smooth muscle of the gastrointestinal tract and the uterine smooth muscle.¹⁷⁻¹⁹ Endothelin has
10 also been shown to promote the growth of rat vascular smooth muscle cells which would suggest a possible relevance to arterial hypertrophy.²⁰

 Endothelin receptors are present in high concentration in the peripheral tissues and also in the central nervous system, and cerebral
15 administration of endothelin has been shown to induce behavioral changes in animals, suggesting that endothelin may play an important role in controlling neural functions.²¹

 Endotoxin has been shown to promote the release of endothelin. This finding has suggested that endothelin is an important
20 mediator for endotoxin-induced diseases.²²⁻²³

 A study has shown that cyclosporin added to a renal cell culture, increased endothelin secretion.²⁴ Another study has shown that administration of cyclosporin to rats led to a decrease in the glomerular filtration rate and an increase in the blood pressure, in association with a
25 remarkable increase in the circulating endothelin level. This cyclosporin-induced renal failure can be suppressed by the administration of anti-endothelin antibody.²⁵ These studies suggest that endothelin is significantly involved in the pathogenesis of cyclosporin-induced renal disease.

30 A recent study in patients with congestive heart failure demonstrated a good correlation between the elevated levels of endothelin in the plasma and the severity of the disease.²⁶

 Substances which specifically inhibit the binding of endothelin to its receptor are believed to block the pathophysiological

- 3 -

effects of endothelin and would be a useful method of treatment for a person in need of such therapy. The present invention discloses potent non-peptidic endothelin antagonists.

5 Endothelin is an endogenous substance which directly or indirectly (through the controlled release of various other endogenous substances) induces sustained contraction of vascular or non-vascular smooth muscles. Its excess production or excess secretion is believed to be one of the factors responsible for hypertension, pulmonary hypertension, Raynaud's disease, bronchial asthma, acute renal failure, myocardial infarction, angina pectoris, arteriosclerosis, cerebral
10 vasospasm and cerebral infarction.

The novel compounds of the present invention are useful as non-peptidic endothelin antagonists, and have not been disclosed in any issued patents or patent applications. Fujisawa in European Patent
15 Application EP 457,195, Banyu in EP 436,189 and 460,679, and Takeda in Patent Cooperation Treaty International Publication No. WO 91/13089 have applications disclosing linear and cyclic peptidic compounds as endothelin antagonists. Fujisawa has also disclosed anthraquinone derivatives produced by a fermentation process using Streptomyces sp.
20 No. 89009 in EP 405,421.

A Roussel-Uclaf European Patent Application (EP 498,723) disclosed a series of substituted (1,4-quinolinoxy)methylbiphenyl-carboxylic acids as both endothelin antagonists and angiotensin II antagonists. A patent from Hoffmann-La Roche (EP 510,526) has also
25 appeared claiming the endothelin antagonist properties of a series of N-(4-pyrimidinyl)benzenesulfonamides.

30

- 4 -

REFERENCES


- 1 Nature, 332, 411-415 (1988).
- 2 FEBS Letters, 231, 440-444 (1988).
- 5 3 Biochem. Biophys. Res. Commun., 154, 868-875 (1988).
- 4 TiPS, 13, 103-108, March 1992.
- 5 Japan J. Hypertension, 12, 79 (1989).
- 6 J. Vascular Medicine Biology, 2, 207 (1990).
- 7 J. Am. Med. Association, 264, 2868 (1990).
- 10 8 The Lancet, ii, 207 (1990) and The Lancet, ii, 747-748 (1989).
- 9 Japan. Soc. Cereb. Blood Flow & Metabol., 1, 73 (1989).
- 10 J. Clin. Invest., 83, 1762-1767 (1989).
- 11 Biochem. Biophys. Res. Comm., 157, 1164-1168 (1988).
- 12 Biochem. Biophys. Res. Comm., 155, 167-172 (1989).
- 15 13 Proc. Natl. Acad. Sci. USA, 85, 9797-9800 (1989).
- 14 J. Cardiovasc. Pharmacol., 13, 589-592 (1989).
- 15 Japan. J. Hypertension, 12, 76 (1989).
- 16 Neuroscience Letters, 102, 179-184 (1989).
- 17 FEBS Letters, 247, 337-340 (1989).
- 20 18 Eur. J. Pharmacol., 154, 227-228 (1988).
- 19 Biochem. Biophys. Res. Commun., 159, 317-323 (1989).
- 20 Atherosclerosis, 78, 225-228 (1989).
- 21 Neuroscience Letters, 97, 276-279 (1989).
- 22 Biochem. Biophys. Res. Commun., 161, 1220-1227 (1989).
- 25 23 Acta. Physiol. Scand., 137, 317-318 (1989).
- 24 Eur. J. Pharmacol., 180, 191-192 (1990).
- 25 Kidney Int., 37, 1487-1491 (1990).
- 26 Mayo Clinic Proc., 67, 719-724 (1992).

30

- 5 -

DETAILED DESCRIPTION OF THE INVENTION

This invention relates to compounds of the general Formula I useful in this novel method of treatment:



or a pharmaceutically acceptable salt thereof, wherein:

R¹ is:

- 20
- (a) H,
- (b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl
- each of which is unsubstituted or substituted with a substituent selected from the group consisting of:
- 25
- i) aryl as defined below,
- ii) (C₃-C₇)-cycloalkyl,
- iii) Cl, Br, I, F,
- iv) OH,
- v) NH₂,
- vi) NH(C₁-C₄)-alkyl,
- 30
- vii) N[(C₁-C₄)-alkyl]₂,
- viii) NHSO₂R²,
- ix) CF₃,
- x) COOR², or

- 6 -

- xi) SO_2NHR^3 ; and
- (c) aryl, wherein aryl is defined as phenyl or naphthyl unsubstituted or substituted with 1 or 2 substituents selected from the group consisting of:
- i) Br, I, Cl, F,
 - ii) $(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 - iii) $(\text{C}_1\text{-C}_4)\text{-alkoxy}$,
 - iv) NO_2
 - v) CF_3
 - vi) $\text{SO}_2\text{NR}^3\text{R}^3$,
 - vii) $(\text{C}_1\text{-C}_4)\text{-alkylthio}$,
 - viii) hydroxy,
 - ix) amino,
 - x) $(\text{C}_3\text{-C}_7)\text{-cycloalkyl}$,
 - xi) $(\text{C}_3\text{-C}_{10})\text{-alkenyl}$; and
- (d) heteroaryl, wherein heteroaryl is defined as a 5- or 6-membered heteroaromatic moiety selected from the group consisting of thiophene, furan, thiazole, oxazole, pyridine or pyrimidine, which is unsubstituted, mono- or disubstituted with substituents selected from the group consisting of:
- i) Cl, Br, I, or F,
 - ii) OH,
 - iii) SH,
 - iv) NO_2 ,
 - v) $(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 - vi) $(\text{C}_2\text{-C}_4)\text{-alkenyl}$,
 - vii) $(\text{C}_2\text{-C}_4)\text{-alkynyl}$,
 - viii) $(\text{C}_1\text{-C}_4)\text{-alkoxy}$, or
 - ix) CF_3 ,
- (e) $(\text{C}_1\text{-C}_4)\text{-perfluoroalkyl}$,
- (f) $-\text{O}-(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (g) $-\text{S}(\text{O})_n-(\text{C}_1\text{-C}_9)\text{-alkyl}$,
- (h) $-\text{CONR}^3\text{R}^3$, or

- 7 -

(i) $\text{-NR}^3\text{CO-O-(C}_1\text{-C}_4\text{)-alkyl}$; and

n is: 0 to 2; and

5 J is: (a) -C(=M)- , (b) J and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} or (c) J and L are connected together to form a 6-membered aromatic ring containing one nitrogen atom, substituted with R^{5a} , R^{5b} and R^{6b} ; and

10

K is: (a) -C(=M)- , (b) K and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} , or (c) K and L are connected together to form a 6-membered aromatic ring containing one nitrogen atom, substituted on the carbon atoms with R^{5a} , R^{5b} and R^{6b} ; and

15

one of a or b is a double bond in Formula I provided that when J is -C(=M)- then b is a double bond and when K is -C(=M)- then a is a double bond.

20

L is the point of attachment of the 6-membered fused aromatic ring optionally containing one nitrogen atom; and

M is: O, S or NR^{11} ; and

25

R^2 is:

- (a) H, or
- (b) $\text{(C}_1\text{-C}_6\text{)-alkyl}$; and

30

R^3 is:

- (a) R^2 ,
- (b) $\text{CH}_2\text{-aryl}$,
- (c) aryl, or

- 8 -

(d) (C₃-C₇)-cycloalkyl; andR⁴ groups are independently:

- 5 (a) H,
(b) (C₁-C₆)-alkyl or (C₂-C₆)-alkenyl each of which is
unsubstituted or substituted with one of the following
substituents,
10 i) -OH,
ii) -O-(C₁-C₄)-alkyl,
iii) -S(O)_n-(C₁-C₄)-alkyl,
iv) -NR³-(C₁-C₄)-alkyl,
v) -NHR³,
vi) -COOR³,
vii) -CONHR³,
15 ix) -CONR³R¹¹, or
x) (C₃-C₇)-cycloalkyl,
(c) (C₃-C₇)-cycloalkyl,
(d) F, Cl, Br, I,
(e) CF₃,
20 (f) -COOR³,
(g) -CONR³R¹¹,
(h) -NR³R¹¹,
(i) -NR³CONR³R¹¹,
(j) -NR³COOR¹¹,
25 (k) -SO₂NR³R¹¹,
(l) -O-(C₁-C₄)-alkyl,
(m) -S(O)_n-(C₁-C₄)-alkyl, or
(n) -NHSO₂R¹¹; and

30 R^{5a} and R^{5b} are independently:

- (a) H,
(b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl,
(c) Cl, Br, I, F,
(d) CF₃, or

- 9 -

- (e) when R^{5a} and R^{5b} are bonded to adjacent carbon atoms, they can be joined to form a phenyl ring;

R^{6a} and R^{6b} are independently:

- 5 (a) H,
 (b) aryl-(C_1 - C_4)-alkyl,
 (c) heteroaryl-(C_1 - C_4)-alkyl,
 (d) (C_1 - C_6)-alkyl, which is unsubstituted or substituted
 10 with a substituent selected from the group consisting of: $-\text{CON}(\text{R}^3)_2$, -heteroaryl, $-\text{S}(\text{O})_n-\text{R}^{15}$, -tetrazol-5-yl, $-\text{CONHSO}_2\text{R}^{15}$, $-\text{SO}_2\text{NH}$ -heteroaryl, $-\text{SO}_2\text{NHCOR}^{15}$, $-\text{PO}(\text{OR}^2)_2$, $-\text{PO}(\text{OR}^3)_2$, $-\text{SO}_2\text{NH-CN}$, $-\text{NR}^2\text{COOR}^{15}$, -OH, $-\text{NH}_2$, guanidino, (C_1 - C_4)-alkoxy, (C_1 - C_4)-alkylthio, (C_1 - C_4)-alkylamino, (C_1 - C_4)-dialkylamino, $-\text{COOR}^3$, $-\text{CONHR}^3$, $-\text{O-COR}^3$, or
 15 aryl,
 (e) $-\text{CO-aryl}$,
 (f) (C_3 - C_7)-cycloalkyl,
 20 (g) Cl, Br, I, F,
 (h) $-\text{OR}^{11}$,
 (i) $-\text{SH}$,
 (j) $-\text{S}(\text{O})_n-(\text{C}_1\text{-}\text{C}_4)\text{-alkyl}$,
 (k) $-\text{COR}^3$,
 (l) $-\text{CO}_2\text{H}$,
 25 (m) $-\text{CO}_2-(\text{C}_1\text{-}\text{C}_4)\text{-alkyl}$,
 (n) $-\text{SO}_3\text{H}$,
 (o) $-\text{NR}^2\text{R}^{15}$,
 (p) $-\text{NR}^2\text{COR}^{15}$,
 (q) $-\text{NR}^2\text{COOR}^{15}$,
 30 (r) $-\text{SO}_2\text{NHR}^3$,
 (s) $-\text{SO}_2\text{NR}^2\text{R}^3$,
 (t) $-\text{NO}_2$,
 (u) $-\text{NHSO}_2\text{CF}_3$,

- 10 -

- 5 (v) $-\text{CONR}^3\text{R}^3$,
(w) $-(\text{C}_1-\text{C}_4)\text{-perfluoroalkyl}$,
(x) $-\text{COOR}^2$,
(y) $-\text{SO}_3\text{H}$,
(z) $-\text{N}(\text{R}^2)\text{SO}_2\text{R}^{15}$,
(aa) $-\text{NR}^2\text{CONR}^3\text{R}^{15}$,
(bb) $-\text{OC}(=\text{O})\text{NR}^{15}\text{R}^3$,
(cc) $-\text{aryl}$, wherein aryl is phenyl or naphthyl, which is
10 unsubstituted or substituted as defined in $\text{R}^1(\text{c})$,
(dd) $-\text{NHSO}_2\text{CF}_3$,
(ee) $-\text{SO}_2\text{NH-heteroaryl}$, wherein heteroaryl is
unsubstituted or substituted as defined in $\text{R}^1(\text{d})$,
(ff) $-\text{SO}_2\text{NHCOR}^{15}$,
(gg) $-\text{CONHSO}_2\text{R}^{15}$,
15 (hh) $-\text{PO}(\text{OR}^2)_2$,
(ii) $-\text{tetrazol-5-yl}$,
(jj) $-\text{CONH}(\text{tetrazol-5-yl})$, or
(kk) $-\text{SO}_2\text{NHCN}$; and

20

 R^7 is:

- (a) H,
(b) $(\text{C}_1-\text{C}_6)\text{-alkyl}$, unsubstituted or substituted with:
25 (i) $-\text{aryl}$,
(ii) $-(\text{C}_3-\text{C}_7)\text{-cycloalkyl}$,
(iii) $-\text{NR}^3\text{R}^{11}$,
(iv) $-\text{morpholin-4-yl}$,
(v) $-\text{OH}$,
(vi) $-\text{CO}_2\text{R}^3$, or
30 (vii) $-\text{CON}(\text{R}^3)_2$,
(c) aryl , unsubstituted or substituted with a substituent
selected from the group consisting of:

- 11 -

- i) (C₁-C₄)-alkyl,
- ii) -O-(C₁-C₄)-alkyl,
- iii) -CONR³R¹¹,
- iv) F, Cl, Br or I, or
- v) -COOR³;

5

R⁸ is:

- (a) H,
- (b) (C₁-C₄)-alkyl, wherein alkyl is unsubstituted or substituted with one or two substituents selected from the group consisting of:
 - i) -OH,
 - ii) -NR³R¹¹,
 - iii) -COOR³,
 - iv) -CONHR³, or
 - v) -CONR³R¹¹;

10

15

R⁹ and R¹⁰ are independently:

- (a) H,
- (b) (C₁-C₆)-alkyl, unsubstituted or substituted with (C₃-C₇)-cycloalkyl,
- (c) (C₂-C₆)-alkenyl,
- (d) (C₂-C₆)-alkynyl,
- (e) Cl, Br, F, I,
- (f) (C₁-C₆)-alkoxy,
- (g) when R⁹ and R¹⁰ are on adjacent carbons, they can be joined to form a phenyl ring,
- (h) perfluoro-(C₁-C₆)-alkyl,
- (i) (C₃-C₇)-cycloalkyl, unsubstituted or substituted with (C₁-C₆)-alkyl,
- (j) aryl,
- (k) (C₁-C₆)-alkyl-S(O)_n-(CH₂)_n-,
- (l) hydroxy-(C₁-C₆)-alkyl or dihydroxy-(C₁-C₆)-alkyl,
- (m) -CF₃,

20

25

30

- 12 -

- 5
- (n) $-\text{CO}_2\text{R}^3$,
 - (o) $-\text{OH}$,
 - (p) $-\text{NR}^3\text{R}^{11}$,
 - (q) $-\text{[(C}_1\text{-C}_6\text{)-alkyl]NR}^3\text{R}^{11}$,
 - (r) $-\text{NO}_2$,
 - (s) $-(\text{CH}_2)_n\text{-SO}_2\text{-N(R}^3)_2$,
 - (t) $-\text{NR}^3\text{CO-(C}_1\text{-C}_4\text{)-alkyl}$, or
 - (u) $-\text{CON(R}^3)_2$;

10

R¹¹ is:

- 15
- (a) H,
 - (b) $(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 - (c) allyl,
 - (d) $(\text{C}_3\text{-C}_6)\text{-cycloalkyl}$,
 - (e) $(\text{C}_1\text{-C}_4)\text{-acyl}$,
 - (f) benzyl, or
 - (g) phenyl; and

20

R¹², R¹³, R^{14a} and R^{14b} are independently:

- 25
- (a) H,
 - (b) F, Cl, Br, or I,
 - (c) $-\text{NO}_2$,
 - (d) $-\text{NH}_2$,
 - (e) $-\text{NH(C}_1\text{-C}_4\text{)-alkyl}$,
 - (f) $-\text{N[(C}_1\text{-C}_4\text{)-alkyl]}_2$,
 - (g) $-\text{SO}_2\text{NHR}^3$,
 - (h) $-\text{CF}_3$,
 - (i) $(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 - (j) $-\text{OR}^3$,
 - (k) $-\text{S(O)}_n\text{-(C}_1\text{-C}_4\text{)-alkyl}$,
 - (l) $-\text{NHCO-(C}_1\text{-C}_4\text{)-alkyl}$,
 - (m) $-\text{NHCO-O(C}_1\text{-C}_4\text{)-alkyl}$,
 - (n) $-\text{CH}_2\text{O-(C}_1\text{-C}_4\text{)-alkyl}$,
- 30

- 13 -

- (o) $-\text{O}-(\text{CH}_2)_m-\text{OR}^3$,
- (p) $-\text{CONR}^3\text{R}^{11}$, or
- (q) $-\text{COOR}^3$ and

5 m is 2, 3, or 4; and

R^{12} and R^{13} on adjacent carbon atoms can be joined together to form a ring structure:

10



A represents:

15

a) $-\text{Y}-\text{C}(\text{R}^4)=\text{C}(\text{R}^4)-$,

b) $-\text{Y}-\text{C}(\text{R}^4)=\text{N}-$,

c) $-\text{Y}-\text{N}=\text{C}(\text{R}^4)-$,

d) $-\text{Y}-[\text{C}(\text{R}^8)(\text{R}^8)]_s-\text{Y}-$,

20

e) $-\text{Y}-\text{C}(\text{R}^8)(\text{R}^8)-\text{C}(\text{R}^8)(\text{R}^8)-$,

f) $-\text{C}(\text{R}^4)=\text{C}(\text{R}^4)-\text{Y}-$,

g) $-\text{N}=\text{C}(\text{R}^4)-\text{Y}-$,

h) $-\text{C}(\text{R}^8)(\text{R}^8)-\text{C}(\text{R}^8)(\text{R}^8)-\text{Y}-$, or

25

i) $-\text{C}(\text{R}^4)=\text{C}(\text{R}^4)-\text{C}(\text{R}^4)=\text{C}(\text{R}^4)-$; and

s is 1 or 2; and

30

Y is $-\text{O}-$, $-\text{S}(\text{O})_n-$ and NR^3 ; and

X is:

(a) $-\text{O}-$,

(b) $-\text{S}(\text{O})_n-$,

- 14 -

- 5
- (c) $-\text{NR}^3-$
 - (d) $-\text{CH}_2\text{O}-$,
 - (e) $-\text{CH}_2\text{S}(\text{O})_n-$,
 - (f) $-\text{CH}_2\text{NR}^3-$,
 - (g) $-\text{OCH}_2-$,
 - (h) $-\text{NR}^3\text{CH}_2-$,
 - (i) $-\text{S}(\text{O})_n\text{CH}_2-$, or
 - (j) single bond; and

10 R^{15} is:

- (a) aryl, or
- (b) (C_1-C_4) -alkyl, is unsubstituted or substituted with:
 - i) NH_2 ,
 - ii) $\text{NH}[(\text{C}_1-\text{C}_4)\text{-alkyl}]$,
 - 15 iii) $\text{N}[(\text{C}_1-\text{C}_4)\text{-alkyl}]_2$,
 - iv) CO_2H ,
 - v) $\text{CO}_2(\text{C}_1-\text{C}_4)\text{-alkyl}$,
 - vi) OH ,
 - vii) SO_3H , or
 - 20 viii) SO_2NH_2 ; and


Z is:

- (a) $-\text{CO}_2\text{H}$,
- (b) $-\text{CO}_2\text{R}^{16}$,
- 25 (c) $-\text{tetrazol-5-yl}$,
- (d) $-\text{CONH}(\text{tetrazol-5-yl})$
- (e) $-\text{CONHSO}_2\text{-aryl}$, wherein aryl is phenyl or naphthyl,
which is unsubstituted or substituted as defined in
 $\text{R}^1(\text{c})$,
- 30 (f) $-\text{CONHSO}_2-(\text{C}_1-\text{C}_8)\text{-alkyl}$, wherein the alkyl group is
unsubstituted or substituted with a substituent selected
from the group consisting of: $-\text{OH}$, $-\text{SH}$, $-\text{O}(\text{C}_1-\text{C}_4)\text{-}$
alkyl, $-\text{S}-(\text{C}_1-\text{C}_4)\text{-alkyl}$, $-\text{CF}_3$, Cl , Br , F , I , $-\text{NO}_2$,

- 15 -

- CO₂H, -CO₂-(C₁-C₄)-alkyl, -NH₂, -NH[(C₁-C₄)-alkyl], -N[(C₁-C₄)-alkyl]₂; and
- (g) -CONHSO₂-(C₁-C₄)-perfluoroalkyl,
- (h) -CONHSO₂-heteroaryl, wherein heteroaryl is
 5 unsubstituted or substituted as defined in R¹(d),
- (i) -CONHSO₂NR³R³,
- (j) -SO₂NHCO-aryl, wherein aryl is phenyl or naphthyl,
 which is unsubstituted or substituted as defined in
 R¹(c),
- 10 (k) -SO₂NHCO-(C₁-C₈)-alkyl, wherein the alkyl group is
 unsubstituted or substituted with a substituent selected
 from the group consisting of: -OH, -SH, -O(C₁-C₄)-
 alkyl, -S-(C₁-C₄)-alkyl, -CF₃, Cl, Br, F, I, -NO₂,
 -CO₂H, -CO₂-(C₁-C₄)-alkyl, -NH₂, -NH[(C₁-C₄)-
 15 alkyl], -N[(C₁-C₄)-alkyl]₂,
- (l) -SO₂NHCO-(C₁-C₄)-perfluoroalkyl,
- (m) -SO₂NHCO-heteroaryl, wherein heteroaryl is
 unsubstituted or substituted as defined in R¹(d),
- (n) -SO₂NHCONR³R³,
- 20 (o) -PO(OH)₂,
- (p) -PO(OR²)₂, or
- (q) -PO(OH)(OR²); and
- R¹⁶ is:
- 25 (a) (C₁-C₄)-alkyl,
- (b) CHR¹⁷-O-COR¹⁸,
- (c) CH₂CH₂-N[(C₁-C₂)-alkyl]₂,
- (d) CH₂CH₂-N[CH₂CH₂]₂O,
- (e) (CH₂CH₂O)_y-O-[(C₁-C₄)-alkyl], wherein y is 1 or 2,
- 30 (f) aryl or CH₂-aryl, where aryl is as defined above or
 optionally substituted with CO₂(C₁-C₄)-alkyl,

- 16 -



R¹⁷ and R¹⁸ independently are: (C₁-C₆)-alkyl or phenyl.

An embodiment of the invention is when:

R¹ is:

- (a) H,
- (b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl
each of which is unsubstituted or substituted with a
substituent selected from the group consisting of:
 - i) (C₃-C₇)-cycloalkyl,
 - ii) CF₃,
 - iii) (C₁-C₄)-alkylthio,
 - iv) (C₁-C₄)-alkoxy,
- (c) (C₁-C₄)-perfluoroalkyl,
- (d) -O-(C₁-C₆)-alkyl,
- (e) -S(O)_n-(C₁-C₉)-alkyl,
- (f) -CONR³R³, or
- (g) -NR³CO-O-(C₁-C₄)-alkyl; and

- 17 -

n is: 0, 1, or 2; and

J is: (a) $-C(=M)-$, (b) J and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} or (c) J and L are connected together to form a 6-membered aromatic ring containing one nitrogen atom, substituted with R^{5a} , R^{5b} and R^{6b} ; and

K is: (a) $-C(=M)-$, or (b) K and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} , or (c) K and L are connected together to form a six-membered aromatic ring containing one nitrogen atom substituted with R^{5a} , R^{5b} and R^{6b} provided that one and only one of J and K is $-C(=M)-$; and

one of a or b is a double bond in Formula I provided that when J is $-C(=M)-$, then b is a double bond and when K is $-C(=M)-$, then a is a double bond;

L is: the point of attachment of the 6-membered fused aromatic ring optionally containing one nitrogen atom; and

M is: O, S or NR^{11} ; and

R^2 is:

- (a) H,
- (b) (C_1-C_6) -alkyl; and

R^3 is:

- (a) R^2 ,
- (b) $-CH_2$ -aryl, or
- (c) aryl; and

- 18 -

R⁴ groups are independently:

- 5
- (a) H,
- (b) (C₁-C₆)-alkyl or (C₂-C₆)-alkenyl each of which is unsubstituted or substituted with one of the following substituents,
- 10
- i) -OH,
- ii) -O-(C₁-C₄)-alkyl,
- iii) -S(O)_n-(C₁-C₄)-alkyl,
- iv) -NR³-(C₁-C₄)-alkyl,
- v) -NHR³,
- vi) -COOR³,
- vii) -CONHR³,
- ix) -CONR³R¹¹, or
- x) (C₃-C₇)-cycloalkyl,
- 15
- (c) (C₃-C₇)-cycloalkyl,
- (d) F, Cl, Br, I,
- (e) CF₃,
- (f) -COOR³,
- (g) -CONR³R¹¹,
- 20
- (h) -NR³R¹¹,
- (i) -NR³CONR³R¹¹,
- (j) -NR³COOR¹¹,
- (k) -SO₂NR³R¹¹,
- (l) -O-(C₁-C₄)-alkyl,
- 25
- (m) -S(O)_n-(C₁-C₄)-alkyl, or
- (n) -NHSO₂R¹¹; and

R^{5a} and R^{5b} are independently:

- 30
- (a) H,
- (b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl,
- (c) Cl, Br, I, F,
- (d) CF₃, or
- (e) when R^{5a} and R^{5b} are bonded to adjacent carbon atoms, they can be joined to form a phenyl ring;

- 19 -

R^{6a} and R^{6b} are independently:

- (a) H,
- (b) aryl-(C₁-C₄)-alkyl,
- (c) heteroaryl-(C₁-C₄)-alkyl,
- (d) (C₁-C₆)-alkyl, is unsubstituted or substituted with a substituent selected from the group consisting of:
 -CON(R³)₂, -heteroaryl, -S(O)_n-R¹⁵, -tetrazol-5-yl,
 -CONHSO₂R¹⁵, -SO₂NH-heteroaryl,
 -SO₂NHCOR¹⁵, -PO(OR²)₂, -PO(OR³)₂, -SO₂NH-CN,
 -NR²COOR¹⁵, -OH, -NH₂, guanidino, (C₁-C₄)-alkoxy,
 (C₁-C₄)-alkylthio, (C₁-C₄)-alkylamino, (C₁-C₄)-dialkylamino,
 -COOR³, -CONHR³, -O-COR³, or aryl,
- (e) -CO-aryl,
- (f) (C₃-C₇)-cycloalkyl,
- (g) Cl, Br, I, F,
- (h) -OR¹¹,
- (i) -SH,
- (j) -S(O)_n-(C₁-C₄)-alkyl,
- (k) -COR³,
- (l) -CO₂H,
- (m) -CO₂-(C₁-C₄)-alkyl,
- (n) -SO₃H,
- (o) -NR²R¹⁵,
- (p) -NR²COR¹⁵,
- (q) -NR²COOR¹⁵,
- (r) -SO₂NHR³,
- (s) -SO₂NR²R³,
- (t) -NO₂,
- (u) -NHSO₂CF₃,
- (v) -CONR³R³,
- (w) -(C₁-C₄)-perfluoroalkyl,

- 20 -

- 5
- (x) $-\text{COOR}^2$,
 - (y) $-\text{SO}_3\text{H}$,
 - (z) $-\text{N}(\text{R}^2)\text{SO}_2\text{R}^{15}$,
 - (aa) $-\text{NR}^2\text{CONR}^3\text{R}^{15}$,
 - (bb) $-\text{OC}(=\text{O})\text{NR}^{15}\text{R}^3$,
 - (cc) -aryl, wherein aryl is phenyl or naphthyl, which is unsubstituted or substituted as defined in $\text{R}^1(\text{c})$,
 - (dd) $-\text{NHSO}_2\text{CF}_3$,
 - (ee) $-\text{SO}_2\text{NH}$ -heteroaryl, wherein heteroaryl is
- 10
- unsubstituted or substituted as defined in $\text{R}^1(\text{d})$,
 - (ff) $-\text{SO}_2\text{NHCOR}^{15}$,
 - (gg) $-\text{CONHSO}_2\text{R}^{15}$,
 - (hh) $-\text{PO}(\text{OR}^2)_2$,
 - (ii) -tetrazol-5-yl,
 - (jj) $-\text{CONH}(\text{tetrazol-5-yl})$, or
 - (kk) $-\text{SO}_2\text{NHCN}$; and
- 15

 R^7 is:

- 20
- (a) H,
 - (b) $(\text{C}_1\text{-C}_6)$ -alkyl, unsubstituted or substituted with:
 - (i) -aryl,
 - (ii) $-(\text{C}_3\text{-C}_7)$ -cycloalkyl,
 - (iii) $-\text{NR}^3\text{R}^{11}$,
 - (iv) -morpholin-4-yl,
 - (v) -OH,
 - (vi) $-\text{CO}_2\text{R}^3$, or
 - (vii) $-\text{CON}(\text{R}^3)_2$,
 - (c) aryl, unsubstituted or substituted with a substituent
- 25
- 30
- selected from the group consisting of:
 - i) $(\text{C}_1\text{-C}_4)$ -alkyl,
 - ii) $-\text{O}(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 - iii) $-\text{CONR}^3\text{R}^{11}$,
 - iv) F, Cl, Br or I, or
 - v) $-\text{COOR}^3$;

- 21 -

R^9 and R^{10} are independently:

- (a) H,
- (b) (C_1-C_6) -alkyl, unsubstituted or substituted with
 (C_3-C_7) -cycloalkyl,
- (c) (C_2-C_6) -alkenyl,
- (d) (C_2-C_6) -alkynyl,
- (e) Cl, Br, F, I,
- (f) (C_1-C_6) -alkoxy,
- (g) when R^9 and R^{10} are on adjacent carbons, they can
 be joined to form a phenyl ring,
- (h) perfluoro- (C_1-C_6) -alkyl,
- (i) (C_3-C_7) -cycloalkyl, unsubstituted or substituted with
 (C_1-C_6) -alkyl,
- (j) aryl,
- (k) (C_1-C_6) -alkyl-S(O)_n-(CH₂)_n-,
- (l) hydroxy- (C_1-C_6) -alkyl or dihydroxy- (C_1-C_6) -alkyl,
- (m) -CF₃,
- (n) -CO₂R³,
- (o) -OH,
- (p) -NR³R¹¹,
- (q) -[(C₁-C₆)-alkyl]NR³R¹¹,
- (r) -NO₂,
- (s) -(CH₂)_n-SO₂-N(R³)₂,
- (t) -NR³CO-(C₁-C₄)-alkyl, or
- (u) -CON(R³)₂;

R^{12} , R^{13} , R^{14a} and R^{14b} are independently:

- (a) H,
- (b) F, Cl, Br, or I,
- (c) -NO₂,
- (d) -NH₂,
- (e) -NH(C₁-C₄)-alkyl,
- (f) -N[(C₁-C₄)-alkyl]₂,

- 22 -

- (g) $-\text{SO}_2\text{NHR}^3$,
 (h) $-\text{CF}_3$,
 (i) $(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 (j) $-\text{OR}^3$,
 5 (k) $-\text{S}(\text{O})_n(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 (l) $-\text{NHCO}(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 (m) $-\text{NHCO-O}(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 (n) $-\text{CH}_2\text{O}(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 (o) $-\text{O}(\text{CH}_2)_m\text{-OR}^3$,
 10 (p) $-\text{CONR}^3\text{R}^{11}$, or
 (q) $-\text{COOR}^3$ and

m is 2, 3, or 4; and

- 15 R^{12} and R^{13} on adjacent carbon atoms can be joined together to
 form a ring structure:



20

A represents:

- a) $-\text{Y-C}(\text{R}^4)=\text{C}(\text{R}^4)-$,
 b) $-\text{Y-C}(\text{R}^4)=\text{N}-$,
 25 c) $-\text{Y-N}=\text{C}(\text{R}^4)-$,
 d) $-\text{Y}-[\text{C}(\text{R}^8)(\text{R}^8)]_s-\text{Y}-$,
 e) $-\text{Y-C}(\text{R}^8)(\text{R}^8)-\text{C}(\text{R}^8)(\text{R}^8)-$,
 f) $-\text{C}(\text{R}^4)=\text{C}(\text{R}^4)-\text{Y}-$,
 30 g) $-\text{N}=\text{C}(\text{R}^4)-\text{Y}-$,
 h) $-\text{C}(\text{R}^8)(\text{R}^8)-\text{C}(\text{R}^8)(\text{R}^8)-\text{Y}-$, or
 i) $-\text{C}(\text{R}^4)=\text{C}(\text{R}^4)-\text{C}(\text{R}^4)=\text{C}(\text{R}^4)-$; and

- 23 -

s is 1 or 2; and

Y is -O-, -S(O)_n- and NR³; and

5

X is:

- (a) -O-,
- (b) -S(O)_n-,
- (c) -NR³-,
- (d) -CH₂O-,
- (e) -CH₂S(O)_n-,
- (f) -CH₂NR³-,
- (g) -OCH₂-,
- (h) -NR³CH₂-,
- (i) -S(O)_nCH₂-, or
- (j) single bond; and

10

15

R¹⁵ is:

- (a) aryl, or
- (b) (C₁-C₄)-alkyl, is unsubstituted or substituted with:
 - i) NH₂,
 - ii) NH[(C₁-C₄)-alkyl],
 - iii) N[(C₁-C₄)-alkyl]₂,
 - iv) CO₂H,
 - v) CO₂(C₁-C₄)-alkyl,
 - vi) OH,
 - vii) SO₃H, or
 - viii) SO₂NH₂; and

20

25

30

Z is:

- (a) -CO₂H,
- (b) -CO₂R¹⁶,
- (c) -tetrazol-5-yl,
- (d) -CONH(tetrazol-5-yl),

- 24 -





- (e) -CONHSO₂-aryl, wherein aryl is phenyl or naphthyl, which is unsubstituted or substituted as defined in R¹(c),
- (f) -CONHSO₂-(C₁-C₈)-alkyl, wherein the alkyl group is unsubstituted or substituted with a substituent selected from the group consisting of: -OH, -SH, -O(C₁-C₄)-alkyl, -S-(C₁-C₄)-alkyl, -CF₃, Cl, Br, F, I, -NO₂, -CO₂H, -CO₂-(C₁-C₄)-alkyl, -NH₂, -NH[(C₁-C₄)-alkyl], -N[(C₁-C₄)-alkyl]₂; and
- (g) -CONHSO₂-(C₁-C₄)-perfluoroalkyl,
- (h) -CONHSO₂-heteroaryl, wherein heteroaryl is unsubstituted or substituted as defined in R¹(d),
- (i) -CONHSO₂NR³R³,
- (j) -SO₂NHCO-aryl, wherein aryl is phenyl or naphthyl, which is unsubstituted or substituted as defined in R¹(c),
- (k) -SO₂NHCO-(C₁-C₈)-alkyl, wherein the alkyl group is unsubstituted or substituted with a substituent selected from the group consisting of: -OH, -SH, -O(C₁-C₄)-alkyl, -S-(C₁-C₄)-alkyl, -CF₃, Cl, Br, F, I, -NO₂, -CO₂H, -CO₂-(C₁-C₄)-alkyl, -NH₂, -NH[(C₁-C₄)-alkyl], -N[(C₁-C₄)-alkyl]₂; and
- (l) -SO₂NHCO-(C₁-C₄)-perfluoroalkyl,
- (m) -SO₂NHCO-heteroaryl, wherein heteroaryl is unsubstituted or substituted as defined in R¹(d), or
- (n) -SO₂NHCONR³R³; and

R¹⁶ is:

- (a) (C₁-C₄)-alkyl,
- (b) CHR¹⁷-O-COR¹⁸,
- (c) CH₂CH₂-N[(C₁-C₂)-alkyl]₂,
- (d) CH₂CH₂-N[CH₂CH₂]₂O,
- (e) (CH₂CH₂O)_y-O-[(C₁-C₄)-alkyl], wherein y is 1 or 2,

- 25 -

- (f) aryl or CH_2 -aryl, where aryl is as defined above or optionally substituted with $\text{CO}_2(\text{C}_1\text{-C}_4)\text{-alkyl}$,



R^{17} and R^{18} independently are: $(\text{C}_1\text{-C}_6)\text{-alkyl}$ or phenyl.

A class of this embodiment of the invention is when:

R^1 is:

- (a) H,
 (b) $(\text{C}_1\text{-C}_6)\text{-alkyl}$, $(\text{C}_2\text{-C}_6)\text{-alkenyl}$ or $(\text{C}_2\text{-C}_6)\text{-alkynyl}$
 each of which is unsubstituted or substituted with a
 substituent selected from the group consisting of:
 i) $(\text{C}_3\text{-C}_7)\text{-cycloalkyl}$,
 ii) CF_3 ,
 iii) $(\text{C}_1\text{-C}_4)\text{-alkylthio}$,
 iv) $(\text{C}_1\text{-C}_4)\text{-alkoxy}$,
 (c) $(\text{C}_1\text{-C}_4)\text{-perfluoroalkyl}$,
 (d) $\text{-CONR}^3\text{R}^3$, or

- 26 -

(e) $-\text{NR}^3\text{CO}-\text{O}-(\text{C}_1-\text{C}_4)\text{-alkyl}$; and

n is: 0, 1, or 2; and

5 J is: (a) $-\text{C}(=\text{M})-$, or (b) J and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} ;

10 K is: (a) $-\text{C}(=\text{M})-$, or (b) K and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} , provided that one and only one of J and K is $-\text{C}(=\text{M})-$; and

15 one of a or b is a double bond in Formula I provided that when J is $-\text{C}(=\text{M})-$, then b is a double bond and when K is $-\text{C}(=\text{M})-$, then a is a double bond;

L is: the point of attachment of the 6-membered fused aromatic ring optionally containing one nitrogen atom; and

20

M is: O, S or NR^{11} ; and

R^2 is:

25 (a) H,
(b) $(\text{C}_1-\text{C}_6)\text{-alkyl}$; and

R^3 is:

30 (a) R^2 ,
(b) $-\text{CH}_2\text{-aryl}$, or
(c) aryl; and

R^4 groups are independently:

(a) H,

- 27 -

(b) (C₁-C₆)-alkyl or (C₂-C₆)-alkenyl each of which is unsubstituted or substituted with one of the following substituents,

- i) -OH,
- ii) -O-(C₁-C₄)-alkyl,
- iii) -S(O)_n-(C₁-C₄)-alkyl,
- iv) -NR³-(C₁-C₄)-alkyl,
- v) -NHR³,
- vi) -COOR³,
- vii) -CONHR³,
- ix) -CONR³R¹¹, or
- x) (C₃-C₇)-cycloalkyl,

(c) (C₃-C₇)-cycloalkyl,

(d) F, Cl, Br, I,

(e) CF₃,

(f) -COOR³,

(g) -CONR³R¹¹,

(h) -NR³R¹¹,

(i) -NR³CONR³R¹¹,

(j) -NR³COOR¹¹,

(k) -SO₂NR³R¹¹,

(l) -O-(C₁-C₄)-alkyl,

(m) -S(O)_n-(C₁-C₄)-alkyl, or

(n) -NHSO₂R¹¹; and

R^{5a} and R^{5b} are independently:

(a) H,

(b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl,

(c) Cl, Br, I, F,

(d) CF₃, or

(e) when R^{5a} and R^{5b} are bonded to adjacent carbon atoms, they can be joined to form a phenyl ring;

R^{6a} and R^{6b} are independently:

- 28 -

- (a) H,
 (b) aryl-(C₁-C₄)-alkyl,
 (c) heteroaryl-(C₁-C₄)-alkyl,
 (d) (C₁-C₆)-alkyl, is unsubstituted or substituted with a
 5 substituent selected from the group consisting of:
 -CON(R³)₂, -heteroaryl, -S(O)_n-R¹⁵, -tetrazol-5-yl,
 -CONHSO₂R¹⁵, -SO₂NH-heteroaryl,
 -SO₂NHCOR¹⁵, -PO(OR²)₂, -PO(OR³)₂, -SO₂NH-
 10 CN, -NR²COOR¹⁵, -OH, -NH₂, guanidino, (C₁-C₄)-
 alkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylamino, (C₁-
 C₄)-dialkylamino, -COOR³, -CONHR³, -O-COR³, or
 aryl,
 (e) -CO-aryl,
 15 (f) (C₃-C₇)-cycloalkyl,
 (g) Cl, Br, I, F,
 (h) -OR¹¹,
 (i) -SH,
 (j) -S(O)_n-(C₁-C₄)-alkyl,
 (k) -COR³,
 20 (l) -CO₂H,
 (m) -CO₂-(C₁-C₄)-alkyl,
 (n) -SO₃H,
 (o) -NR²R¹⁵,
 (p) -NR²COR¹⁵,
 25 (q) -NR²COOR¹⁵,
 (r) -SO₂NHR³,
 (s) -SO₂NR²R³,
 (t) -NO₂,
 (u) -NHSO₂CF₃,
 30 (v) -CONR³R³,
 (w) -(C₁-C₄)-perfluoroalkyl,
 (x) -COOR²,
 (y) -SO₃H,

- 29 -

- 5
- (z) $-N(R^2)SO_2R^{15}$,
 - (aa) $-NR^2CONR^3R^{15}$,
 - (bb) $-OC(=O)NR^{15}R^3$,
 - (cc) -aryl, wherein aryl is phenyl or naphthyl, which is
unsubstituted or substituted as defined in $R^1(c)$,
 - (dd) $-NHSO_2CF_3$,
 - (ee) $-SO_2NH$ -heteroaryl, wherein heteroaryl is
unsubstituted or substituted as defined in $R^1(d)$,
 - (ff) $-SO_2NHCOR^{15}$,
 - 10 (gg) $-CONHSO_2R^{15}$,
 - (hh) $-PO(OR^2)_2$,
 - (ii) -tetrazol-5-yl,
 - (jj) $-CONH(tetrazol-5-yl)$, or
 - 15 (kk) $-SO_2NHCN$; and

 R^7 is:

- 20
- (a) H,
 - (b) (C_1-C_6) -alkyl, unsubstituted or substituted with:
 - (i) -aryl,
 - (ii) $-(C_3-C_7)$ -cycloalkyl,
 - (iii) $-NR^3R^{11}$,
 - (iv) -morpholin-4-yl,
 - (v) -OH,
 - 25 (vi) $-CO_2R^3$, or
 - (vii) $-CON(R^3)_2$,
 - (c) aryl, unsubstituted or substituted with a substituent
selected from the group consisting of:
 - 30 i) (C_1-C_4) -alkyl,
 - ii) $-O-(C_1-C_4)$ -alkyl,
 - iii) $-CONR^3R^{11}$,
 - iv) F, Cl, Br or I, or
 - v) $-COOR^3$;

- 30 -

R⁹ and R¹⁰ are independently:

- (a) H,
- (b) (C₁-C₆)-alkyl, unsubstituted or substituted with (C₃-C₇)-cycloalkyl,
- 5 (c) Cl, Br, F, I,
- (d) (C₁-C₆)-alkoxy, or
- (e) hydroxy-(C₁-C₆)-alkyl or dihydroxy-(C₁-C₆)-alkyl;

R¹², R¹³, R^{14a} and R^{14b} are independently:

- 10 (a) H,
- (b) F, Cl, Br, or I,
- (c) -NO₂,
- (d) (C₁-C₄)-alkyl,
- (e) -OR³,
- 15 (f) -S(O)_n-(C₁-C₄)-alkyl,
- (g) -NHCO-(C₁-C₄)-alkyl,
- (h) -NHCO-O(C₁-C₄)-alkyl,
- (i) -O-(CH₂)_m-OR³,
- (j) -CONR³R¹¹, or
- 20 (k) -COOR³ and

m is 2, 3, or 4; and

25 R¹² and R¹³ on adjacent carbon atoms can be joined together to form a ring structure:



30

A represents:

- a) -O-C(R⁴)=C(R⁴)-,
- b) -O-C(R⁴)=N-,
- c) -O-[C(R⁸)(R⁸)]_s-O-,

- 31 -

- d) $-C(R^4)=C(R^4)-O-$,
- e) $-N=C(R^4)-O-$, or
- f) $-C(R^4)=C(R^4)-C(R^4)=C(R^4)-$; and

5

s is 1 or 2; and

X is:

10

- (a) $-O-$,
- (b) $-S(O)_n-$, or
- (c) $-NR^3-$; and

 R^{15} is:

15

- (a) aryl, or
- (b) (C_1-C_4) -alkyl, is unsubstituted or substituted with:
 - i) NH_2 ,
 - ii) $NH[(C_1-C_4)\text{-alkyl}]$,
 - iii) $N[(C_1-C_4)\text{-alkyl}]_2$,
 - iv) CO_2H ,
 - v) $CO_2(C_1-C_4)\text{-alkyl}$,
 - vi) OH ,
 - vii) SO_3H , or
 - viii) SO_2NH_2 ; and

20

25


Z is:

- (a) $-CO_2H$,
- (b) -tetrazol-5-yl ,
- (c) $\text{-CONH(tetrazol-5-yl)}$,
- (d) $\text{-CONHSO}_2\text{-phenyl}$ or $\text{-CONHSO}_2\text{-naphthyl}$, wherein phenyl or naphthyl is unsubstituted or substituted as defined in $R^1(c)$, or
- (e) $\text{-CONHSO}_2\text{-heteroaryl}$, wherein heteroaryl is unsubstituted or substituted as defined in $R^1(d)$.

30

- 32 -

Another class of this embodiment of the invention is a compound of Formula II



wherein:

R¹ is:

- 20
- (a) H,
- (b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl
each of which is unsubstituted or substituted with a
substituent selected from the group consisting of:
- 25
- i) (C₃-C₇)-cycloalkyl,
- ii) CF₃,
- iii) (C₁-C₄)-alkylthio,
- iv) (C₁-C₄)-alkoxy,
- (c) (C₁-C₄)-perfluoroalkyl,
- (d) -CONR³R³, or
- 30
- (e) -NR³CO-O-(C₁-C₄)-alkyl; and

n is: 0, 1, or 2; and

- 33 -

J is: (a) $-C(=M)-$, or (b) J and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} ;

5 K is: (a) $-C(=M)-$, or (b) K and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} , provided that one and only one of J and K is $-C(=M)-$; and

10 one of a or b is a double bond in Formula I provided that when J is $-C(=M)-$, then b is a double bond and when K is $-C(=M)-$, then a is a double bond;

15 L is: the point of attachment of the 6-membered fused aromatic ring optionally containing one nitrogen atom; and

M is: O, S or NR^{11} ; and

R^2 is:

- 20 (a) H,
(b) (C_1-C_6) -alkyl; and

R^3 is:

- 25 (a) R^2 ,
(b) $-CH_2$ -aryl, or
(c) aryl; and

R^4 groups are independently:

- 30 (a) H,
(b) (C_1-C_6) -alkyl or (C_2-C_6) -alkenyl each of which is unsubstituted or substituted with one of the following substituents,
i) $-OH$,
ii) $-O-(C_1-C_4)$ -alkyl,

- 34 -

- 5
10
15
20
- iii) $-S(O)_n-(C_1-C_4)\text{-alkyl}$,
 - iv) $-NR^3-(C_1-C_4)\text{-alkyl}$,
 - v) $-NHR^3$,
 - vi) $-COOR^3$,
 - vii) $-CONHR^3$,
 - ix) $-CONR^3R^{11}$, or
 - x) $(C_3-C_7)\text{-cycloalkyl}$,
 - (c) $(C_3-C_7)\text{-cycloalkyl}$,
 - (d) F, Cl, Br, I,
 - (e) CF_3 ,
 - (f) $-COOR^3$,
 - (g) $-CONR^3R^{11}$,
 - (h) $-NR^3R^{11}$,
 - (i) $-NR^3CONR^3R^{11}$,
 - (j) $-NR^3COOR^{11}$,
 - (k) $-SO_2NR^3R^{11}$,
 - (l) $-O-(C_1-C_4)\text{-alkyl}$,
 - (m) $-S(O)_n-(C_1-C_4)\text{-alkyl}$, or
 - (n) $-NH SO_2R^{11}$; and

20 R^{5a} and R^{5b} are independently:

- 25
- (a) H,
 - (b) $(C_1-C_6)\text{-alkyl}$, $(C_2-C_6)\text{-alkenyl}$ or $(C_2-C_6)\text{-alkynyl}$,
 - (c) Cl, Br, I, F,
 - (d) CF_3 , or
 - (e) when R^{5a} and R^{5b} are bonded to adjacent carbon atoms, they can be joined to form a phenyl ring;

30 R^{6a} and R^{6b} are independently:

- 30
- (a) H,
 - (b) $\text{aryl}-(C_1-C_4)\text{-alkyl}$,
 - (c) $\text{heteroaryl}-(C_1-C_4)\text{-alkyl}$,
 - (d) $(C_1-C_6)\text{-alkyl}$, is unsubstituted or substituted with a substituent selected from the group consisting of:

- 35 -

- 5
-CON(R³)₂, -heteroaryl, -S(O)_n-R¹⁵, -tetrazol-5-yl,
-CONHSO₂R¹⁵, -SO₂NH-heteroaryl,
-SO₂NHCOR¹⁵, -PO(OR²)₂, -PO(OR³)₂, -SO₂NH-
CN, -NR²COOR¹⁵, -OH, -NH₂, guanidino, (C₁-C₄)-
alkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylamino, (C₁-
C₄)-dialkylamino, -COOR³, -CONHR³, -O-COR³, or
aryl,
- 10
(e) -CO-aryl,
(f) (C₃-C₇)-cycloalkyl,
(g) Cl, Br, I, F,
(h) -OR¹¹,
(i) -SH,
(j) -S(O)_n-(C₁-C₄)-alkyl,
- 15
(k) -COR³,
(l) -CO₂H,
(m) -CO₂-(C₁-C₄)-alkyl,
(n) -SO₃H,
(o) -NR²R¹⁵,
- 20
(p) -NR²COR¹⁵,
(q) -NR²COOR¹⁵,
(r) -SO₂NHR³,
(s) -SO₂NR²R³,
(t) -NO₂,
- 25
(u) -NHSO₂CF₃,
(v) -CONR³R³,
(w) -(C₁-C₄)-perfluoroalkyl,
(x) -COOR²,
(y) -SO₃H,
- 30
(z) -N(R²)SO₂R¹⁵,
(aa) -NR²CONR³R¹⁵,
(bb) -OC(=O)NR¹⁵R³,
(cc) -aryl, wherein aryl is phenyl or naphthyl, which is
unsubstituted or substituted as defined in R¹(c),

- 36 -

- (dd) $-\text{NHSO}_2\text{CF}_3$,
(ee) $-\text{SO}_2\text{NH}$ -heteroaryl, wherein heteroaryl is
unsubstituted or substituted as defined in $\text{R}^1(\text{d})$,
(ff) $-\text{SO}_2\text{NHCOR}^{15}$,
(gg) $-\text{CONHSO}_2\text{R}^{15}$,
(hh) $-\text{PO}(\text{OR}^2)_2$,
(ii) -tetrazol-5-yl,
(jj) $-\text{CONH}(\text{tetrazol-5-yl})$, or
(kk) $-\text{SO}_2\text{NHCN}$; and

 R^7 is:

- (a) H,
(b) $(\text{C}_1\text{-C}_6)$ -alkyl, unsubstituted or substituted with:
(i) -aryl,
(ii) $-(\text{C}_3\text{-C}_7)$ -cycloalkyl,
(iii) $-\text{NR}^3\text{R}^{11}$,
(iv) -morpholin-4-yl,
(v) -OH,
(vi) $-\text{CO}_2\text{R}^3$, or
(vii) $-\text{CON}(\text{R}^3)_2$,
(c) aryl, unsubstituted or substituted with a substituent
selected from the group consisting of:
i) $(\text{C}_1\text{-C}_4)$ -alkyl,
ii) $-\text{O}-(\text{C}_1\text{-C}_4)$ -alkyl,
iii) $-\text{CONR}^3\text{R}^{11}$,
iv) F, Cl, Br or I, or
v) $-\text{COOR}^3$;

 R^9 and R^{10} are independently:

- (a) H,
(b) $(\text{C}_1\text{-C}_6)$ -alkyl, unsubstituted or substituted with
 $(\text{C}_3\text{-C}_7)$ -cycloalkyl,
(c) Cl, Br, F, I,

- 37 -

- (d) (C₁-C₆)-alkoxy, or
- (e) hydroxy-(C₁-C₆)-alkyl or dihydroxy-(C₁-C₆)-alkyl;

R₁₂, R₁₃, R_{14a} and R_{14b} are independently:

- 5 (a) H,
- (b) F, Cl, Br, or I,
- (c) -NO₂,
- (d) (C₁-C₄)-alkyl,
- (e) -OR³,
- 10 (f) -S(O)_n-(C₁-C₄)-alkyl,
- (g) -NHCO-(C₁-C₄)-alkyl,
- (h) -NHCO-O(C₁-C₄)-alkyl,
- (i) -O-(CH₂)_m-OR³,
- (j) -CONR³R¹¹, or
- 15 (k) -COOR³ and

m is 2, 3, or 4; and

R₁₂ and R₁₃ on adjacent carbon atoms can be joined together to form a ring structure:



25

A represents:

- a) -O-C(R⁴)=C(R⁴)-,
- b) -O-C(R⁴)=N-,
- c) -O-[C(R⁸)(R⁸)]_s-O-,
- 30 d) -C(R⁴)=C(R⁴)-O-,
- e) -N=C(R⁴)-O-, or
- f) -C(R⁴)=C(R⁴)-C(R⁴)=C(R⁴)-; and

s is 1 or 2; and

- 38 -

X is:

- (a) -O-,
- (b) -S(O)_n-, or
- (c) -NR³-; and

5

R¹⁵ is:

- (a) aryl, or
- (b) (C₁-C₄)-alkyl, is unsubstituted or substituted with:
 - i) NH₂,
 - ii) NH[(C₁-C₄)-alkyl],
 - 10 iii) N[(C₁-C₄)-alkyl]₂,
 - iv) CO₂H,
 - v) CO₂(C₁-C₄)-alkyl,
 - vi) OH,
 - 15 vii) SO₃H, or
 - viii) SO₂NH₂; and

Z is:

- (a) -CO₂H,
- (b) -tetrazol-5-yl,
- (c) -CONH(tetrazol-5-yl),
- (d) -CONHSO₂-phenyl or -CONHSO₂-naphthyl, wherein
phenyl or naphthyl is unsubstituted or substituted as
defined in R¹(c), or
- (e) -CONHSO₂-heteroaryl, wherein heteroaryl is
25 unsubstituted or substituted as defined in R¹(d).

Another class of this embodiment of the invention is a
compound of Formula III:


30

- 39 -

5

10

15

R¹ is:

20

25

30

- (a) H,
- (b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl
each of which is unsubstituted or substituted with a
substituent selected from the group consisting of:
 - i) (C₃-C₇)-cycloalkyl,
 - ii) CF₃,
 - iii) (C₁-C₄)-alkylthio,
 - iv) C₁-C₄-alkoxy,
- (c) (C₁-C₄)-perfluoroalkyl,
- (d) -CONR³R³, or
- (e) -NR³CO-O-(C₁-C₄)-alkyl; and

n is: 0, 1, or 2; and

R² is:

- (a) H,

- 40 -

(b) (C₁-C₆)-alkyl; andR³ is:

- 5
- (a) R²,
 - (b) -CH₂-aryl, or
 - (c) aryl; and

R⁴ groups are independently:

- 10
- (a) H,
 - (b) (C₁-C₆)-alkyl or (C₂-C₆)-alkenyl each of which is unsubstituted or substituted with one of the following substituents,
 - i) -OH,
 - ii) -O-(C₁-C₄)-alkyl,
 - 15 iii) -S(O)_n-(C₁-C₄)-alkyl,
 - iv) -NR³-(C₁-C₄)-alkyl,
 - v) -NHR³,
 - vi) -COOR³,
 - vii) -CONHR³,
 - 20 ix) -CONR³R¹¹, or
 - x) (C₃-C₇)-cycloalkyl,
 - (c) (C₃-C₇)-cycloalkyl,
 - (d) F, Cl, Br, I,
 - (e) CF₃,
 - 25 (f) -COOR³,
 - (g) -CONR³R¹¹,
 - (h) -NR³R¹¹,
 - (i) -NR³CONR³R¹¹,
 - (j) -NR³COOR¹¹,
 - 30 (k) -SO₂NR³R¹¹,
 - (l) -O-(C₁-C₄)-alkyl,
 - (m) -S(O)_n-(C₁-C₄)-alkyl, or
 - (n) -NHSO₂R¹¹; and

- 41 -

R^{5a} is:

- 5
- (a) H,
 - (b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl,
 - (c) Cl, Br, I, F,
 - (d) CF₃, or
 - (e) when R^{5a} and R^{5b} are bonded to adjacent carbon atoms, they can be joined to form a phenyl ring;

R^{6a} is:

- 10
- (a) H,
 - (b) aryl-(C₁-C₄)-alkyl,
 - (c) heteroaryl-(C₁-C₄)-alkyl,
 - (d) (C₁-C₆)-alkyl, is unsubstituted or substituted with a substituent selected from the group consisting of:
- 15
- CON(R³)₂, -heteroaryl, -S(O)_n-R¹⁵, -tetrazol-5-yl,
 - CONHSO₂R¹⁵, -SO₂NH-heteroaryl,
 - SO₂NHCOR¹⁵, -PO(OR²)₂, -PO(OR³)₂, -SO₂NH-
 - CN, -NR²COOR¹⁵, -OH, -NH₂, guanidino, (C₁-C₄)-
 - alkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylamino, (C₁-
 - 20 C₄)-dialkylamino, -COOR³, -CONHR³, -O-COR³, or
 - aryl,
 - (e) -CO-aryl,
 - (f) (C₃-C₇)-cycloalkyl,
 - (g) Cl, Br, I, F,
 - (h) -OR¹¹,
 - (i) -SH,
 - (j) -S(O)_n-(C₁-C₄)-alkyl,
 - (k) -COR³,
 - (l) -CO₂H,
 - (m) -CO₂-(C₁-C₄)-alkyl,
 - (n) -SO₃H,
 - (o) -NR²R¹⁵,
 - (p) -NR²COR¹⁵,
- 25
- 30

- 42 -

- (q) $-\text{NR}^2\text{COOR}^{15}$,
 (r) $-\text{SO}_2\text{NHR}^3$,
 (s) $-\text{SO}_2\text{NR}^2\text{R}^3$,
 (t) $-\text{NO}_2$,
 5 (u) $-\text{NHSO}_2\text{CF}_3$,
 (v) $-\text{CONR}^3\text{R}^3$,
 (w) $-(\text{C}_1-\text{C}_4)\text{-perfluoroalkyl}$,
 (x) $-\text{COOR}^2$,
 (y) $-\text{SO}_3\text{H}$,
 10 (z) $-\text{N}(\text{R}^2)\text{SO}_2\text{R}^{15}$,
 (aa) $-\text{NR}^2\text{CONR}^3\text{R}^{15}$,
 (bb) $-\text{OC}(=\text{O})\text{NR}^{15}\text{R}^3$,
 (cc) $-\text{aryl}$, wherein aryl is phenyl or naphthyl, which is
 unsubstituted or substituted as defined in $\text{R}^1(\text{c})$,
 15 (dd) $-\text{NHSO}_2\text{CF}_3$,
 (ee) $-\text{SO}_2\text{NH-heteroaryl}$, wherein heteroaryl is
 unsubstituted or substituted as defined in $\text{R}^1(\text{d})$,
 (ff) $-\text{SO}_2\text{NHCOR}^{15}$,
 (gg) $-\text{CONHSO}_2\text{R}^{15}$,
 20 (hh) $-\text{PO}(\text{OR}^2)_2$,
 (ii) $-\text{tetrazol-5-yl}$,
 (jj) $-\text{CONH}(\text{tetrazol-5-yl})$, or
 (kk) $-\text{SO}_2\text{NHCN}$; and

25

R^9 and R^{10} are independently:

- (a) H,
 (b) $(\text{C}_1-\text{C}_6)\text{-alkyl}$, unsubstituted or substituted with
 $(\text{C}_3-\text{C}_7)\text{-cycloalkyl}$,
 30 (c) Cl, Br, F, I,
 (d) $(\text{C}_1-\text{C}_6)\text{-alkoxy}$, or
 (e) hydroxy- $(\text{C}_1-\text{C}_6)\text{-alkyl}$ or dihydroxy- $(\text{C}_1-\text{C}_6)\text{-alkyl}$;

R^{12} , R^{13} , R^{14a} and R^{14b} are independently:

- 43 -

- 5
- (a) H,
 - (b) F, Cl, Br, or I,
 - (c) -NO₂,
 - (d) (C₁-C₄)-alkyl,
 - (e) -OR³,
 - (f) -S(O)_n-(C₁-C₄)-alkyl,
 - (g) -NHCO-(C₁-C₄)-alkyl,
 - (h) -NHCO-O(C₁-C₄)-alkyl,
 - (i) -O-(CH₂)_m-OR³,
 - 10 (j) -CONR³R¹¹, or
 - (k) -COOR³ and

m is 2, 3, or 4; and

- 15 R¹² and R¹³ on adjacent carbon atoms can be joined together to form a ring structure:



A represents:

- 25
- a) -O-C(R⁴)=C(R⁴)-,
 - b) -O-C(R⁴)=N-,
 - c) -O-[C(R⁸)(R⁸)]_s-O-,
 - d) -C(R⁴)=C(R⁴)-O-,
 - e) -N=C(R⁴)-O-, or
 - 30 f) -C(R⁴)=C(R⁴)-C(R⁴)=C(R⁴)-; and

s is 1 or 2; and

- 44 -

X is:

- (a) -O-,
- (b) -S(O)_n-, or
- (c) -NR³-; and

5

R¹⁵ is:

- (a) aryl, or
- (b) (C₁-C₄)-alkyl, is unsubstituted or substituted with:
 - i) NH₂,
 - ii) NH[(C₁-C₄)-alkyl],
 - 10 iii) N[(C₁-C₄)-alkyl]₂,
 - iv) CO₂H,
 - v) CO₂(C₁-C₄)-alkyl,
 - vi) OH,
 - 15 vii) SO₃H, or
 - viii) SO₂NH₂; and

Z is:


20

- (a) -CO₂H,
- (b) -tetrazol-5-yl,
- (c) -CONH(tetrazol-5-yl),
- (d) -CONHSO₂-phenyl or -CONHSO₂-naphthyl, wherein
phenyl or naphthyl is unsubstituted or substituted as
defined in R¹(c), or
- 25 (e) -CONHSO₂-heteroaryl, wherein heteroaryl is as
defined in R¹(d).

30

- 45 -

The following Table further exemplifies the scope of the invention wherein the structural formula is:



and the substituents are as defined in the table below:

20	<u>R¹</u>	<u>R^{6a}</u>	<u>R⁹</u>	<u>R¹⁰</u>	<u>R¹², R¹³, R^{14a}, R^{14b}</u>	<u>Z</u>
	Bu	6-Me	H	Allyl	H, H, H, H	COOH
	Bu	6-Me	H	Pr	H, H, H, H	COOH
	Bu	6-Me	H	Cl	H, H, H, H	COOH
	Pr	6-N(Me)COOiBu	OMe	Cl	H, H, H, H	COOH
25	Pr	6-N(Me)COOiBu	H	Pr	H, H, H, H	COOH
	Pr	6-Me	Pr	Pr	2-Br, 5-Br, 3-OMe, 4-OMe	COOH
	H	8-Me	Pr	Pr	2-Br, 5-Br, 3-OMe, 4-OMe	COOH
	Me	8-Me	Me	Me	3-OMe, 5-OMe	COOH
	H	8-Me	H	Pr	3-OMe, 5-OMe	COOH
30	H	8-Me	Pr	Pr	3-OMe, 5-OMe	COOH
	H	8-Me	H	Pr	3-OMe, 5-OMe	CONHSO ₂ Ph

- 46 -

	<u>R¹</u>	<u>R^{6a}</u>	<u>R⁹</u>	<u>R¹⁰</u>	<u>R¹², R¹³, R^{14a}, R^{14b}</u>	<u>Z</u>
	H	8-Me	Pr	Pr	3-OMe, 5-OMe	CONHSO ₂ Me
	H	8-Me	H	Pr	3-OMe, 5-OMe	CONHSO ₂ Ph(4-iPr)
	i-Pr	8-Br	H	Pr	3-OMe, 5-OMe	CONHSO ₂ Ph(4-iPr)
5	H	6-PhCONH	Bu	H	3-OMe	COOH
	Me	8-Me	Pr	H	3-OMe	COOH
	Me	8-Me	Pr	H	3-OMe	CONHSO ₂ Ph(4-iPr)
	Me	8-Me	Pr	Pr	3-OMe	COOH
	H	8-Me	Cl	Cl	3-OMe	COOH
10	Me	8-Me	Br	Br	3-OMe	COOH
	Ph	8-Me	Cl	Cl	3-OMe	COOH
	H	8-Me	Pr	Pr	4-OMe	COOH
	Me	6-Me	Pr	Pr	4-OMe	COOH
	i-Pr	8-Me	Pr	H	4-OMe	COOH
15	Bu	6-Me	Cl	Cl	4-OMe	COOH
	Me	8-Me	Br	Br	4-OMe	COOH
	H	8-Me	Pr	Pr	2-OMe	COOH
	Me	6-Me	Pr	Pr	2-OMe, 3-OMe	COOH
	i-Pr	8-Me	Pr	H	2-OMe	COOH
20	Bu	6-Me	Cl	Cl	2-OMe	COOH
	Me	8-Me	Br	Br	2-OMe	COOH
	H	8-Me	H	Pr	3-OMe, 4-OMe	COOH
	H	8-Me	Pr	Pr	3-OMe, 4-OMe	COOH
	H	8-Me	H	Pr	3-OMe, 4-OMe	CONHSO ₂ Ph
25	H	8-Me	Pr	Pr	3-OMe, 4-OMe	CONHSO ₂ Me
	H	8-Me	H	Pr	3-OMe, 4-OMe	CONHSO ₂ Ph(4-iPr)

- 47 -

	<u>R¹</u>	<u>R^{6a}</u>	<u>R⁹</u>	<u>R¹⁰</u>	<u>R¹², R¹³, R^{14a}, R^{14b}</u>	<u>Z</u>
	i-Pr	8-Br	H	Pr	3-OMe, 4-OMe	CONHSO ₂ Ph(4-iPr)
	H	6-PhCONH Bu	H	H	2-OMe, 3-OMe	COOH
	Me	8-Me	Pr	H	2-OMe, 3-OMe	COOH
5	Me	8-Me	Pr	H	2-OMe, 3-OMe	CONHSO ₂ Ph(4-iPr)
	Me	8-Me	Pr	Pr	2-OMe, 3-OMe	COOH
	H	8-Me	Cl	Cl	2-OMe, 3-OMe	COOH
	Me	8-Me	Br	Br	2-OMe, 3-OMe	COOH
	H	8-Me	H	Pr	3,4-methylenedioxy	COOH
10	H	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
	H	8-Me	H	Pr	3,4-methylenedioxy	CONHSO ₂ Ph
	H	8-Me	Pr	Pr	3,4-methylenedioxy	CONHSO ₂ Me
	i-Pr	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
	i-Pr	8-Br	Pr	Pr	3,4-methylenedioxy	COOH
15	Ph	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
	(2-Et)Pr	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
	H	8-Me	H	Pr	3,4-methylenedioxy	CONHSO ₂ Ph(4-iPr)
	i-Pr	8-Br	H	Pr	3,4-methylenedioxy	CONHSO ₂ Ph(4-iPr)
	H	6-PhCONH Bu	H	H	3,4-methylenedioxy	COOH
20	Me	8-Me	Pr	H	3,4-methylenedioxy	COOH
	Me	8-Me	Pr	H	3,4-methylenedioxy	CONHSO ₂ Ph(4-iPr)
	Me	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
	H	8-Me	Cl	Cl	3,4-methylenedioxy	COOH
	Me	8-Me	Br	Br	3,4-methylenedioxy	COOH
25	H	8-Me	H	Pr	5-Br; 3,4-methylenedioxy	COOH
	H	8-Me	Pr	Pr	5-Br; 3,4-methylenedioxy	COOH
	H	8-Me	H	Pr	5-Br; 3,4-methylenedioxy	CONHSO ₂ Ph(4-iPr).

- 48 -

The alkyl substituents recited above denote straight and branched chain hydrocarbons of the length specified such as methyl, ethyl, isopropyl, isobutyl, neopentyl, isopentyl, etc.

5 The alkenyl and alkynyl substituents denote alkyl groups as described above which are modified so that each contains a carbon to carbon double bond or triple bond, respectively, such as vinyl, allyl and 2-butenyl.

10 Cycloalkyl denotes rings composed of 3 to 8 methylene groups, each of which may be substituted or unsubstituted with other hydrocarbon substituents, and include for example cyclopropyl, cyclopentyl, cyclohexyl and 4-methylcyclohexyl.

The alkoxy substituent represents an alkyl group as described above attached through an oxygen bridge.

15 The heteroaryl substituent recited above represents any 5- or 6-membered aromatic ring containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, for example, pyridyl, furyl, pyrrolyl, thienyl, isothiazolyl, imidazolyl, pyrazinyl, pyrimidyl, pyrazolyl, purinyl, carbazolyl, isoxazolyl, thiazolyl, and oxazolyl.

20 The reaction schemes described below are reasonably general, it will be understood by those skilled in the art of organic synthesis that one or more functional groups present in a given compound of Formula I may render the molecule incompatible with a particular synthetic sequence. In such a case an alternative synthetic route, an altered order of steps, or a strategy of protection and deprotection may be
25 employed. In all cases the particular reaction conditions, including reagents, solvent, temperature and time, should be chosen so that they are consistent with the nature of the functionality present in the molecule.

30 The compounds of Formula I, and more specifically compounds where R⁷ is hydrogen, can be synthesized using the reactions and techniques described for the synthesis of the non-heterocyclic components in the patent application WO91/12001 (Merck & Co.; published on August 22, 1991 under the Patent Cooperation Treaty) and also US Patent 5,177,095 (Merck & Co.; January 5, 1993).

- 49 -

The reaction schemes described below have been generalized for simplicity. It is further to be understood that in the generalized schemes below, unless specified more narrowly in the text, the alkyl and aryl groups represent unfunctionalized or functionalized derivatives as described before. The leaving group Q present in the alkylating agents is either chloro, bromo, iodo, methanesulfonate, p-toluenesulfonate or triflate.

10

15


20

25

30

- 50 -


Scheme 1



- 51 -

More specifically, the compounds of Formulae II (where X is oxygen, sulphur or appropriately substituted nitrogen and R⁷ is H) can be synthesized as outlined in Scheme 1. The substituted compound 1 may be reacted with the alkylating agent 2 in an appropriate solvent such as alcohols (methanol, ethanol, isopropanol and like), dimethylformamide (DMF), dimethylsulfoxide (DMSO), tetrahydrofuran (THF) and acetone in the presence of an alkali metal salt such as alkoxides, carbonates, hydroxides and hydrides, or an organic base such as trialkylamines and alkyl lithiums to provide compound 3. The Z¹ group present in compound 3 may then be further transformed to provide desired compounds of Formula II.

In general, the alkylating agent 2 can be prepared using methods and techniques outlined in US Patent 5,177,095. More specifically, compound 2 (where Z¹ is COOR and Q is Br) can be synthesized from the substituted arylacetic acids 4 as outlined in Scheme 2. Substituted arylacetic acid 4 is converted to the corresponding ester either by refluxing the acid in an appropriate alcohol in the presence of a catalytic amount of conc. sulfuric acid, or using other conventional methods of esterification. The resulting ester is then refluxed in carbon tetrachloride with N-Bromosuccinimide and a catalytic amount of a radical initiator (e.g., AIBN or benzoylperoxide) to provide the 2-bromo-arylacetic acid ester 5.

Scheme 2


Alternatively, the ester 5 may also be prepared from appropriate aryl aldehydes (Scheme 3). The aldehyde 6 can be reacted with trimethylsilyl cyanide and catalytic amounts of KCN and 18-crown-6 to provide the corresponding trimethylsilyl cyanohydrin 7, which upon

- 52 -

further treatment with the gaseous HCl and alcohol can afford the 2-hydroxy ester **8**. The ester **8** is treated with triphenylphosphine and carbon tetrabromide in methylene chloride to give the 2-bromoarylacetate derivatives **5**.

5

Scheme 3



a. TMSCN, Cat. KCN, CH₂Cl₂, 18-Crown-6; b. HCl(g), EtOH;


c. CBr₄, Ph₃P, CH₂Cl₂

Scheme 4 illustrates a typical synthesis of an alkylating agent **12** (where Ar represents substituted indoles). The appropriately substituted cyanoindole **9** (for a general synthesis of substituted indoles refer to, R. K. Brown, *Indoles, Part One*, Ed. W. J. Houlihan, Vol. 25, Chapter II, Wiley-Interscience, New York, 1972) is reduced with DIBALH to provide the corresponding aldehyde, which is then converted into the N-Boc derivative **10**. Reaction of **10** with the trichloromethide anion [generated from KOH and CHCl₃; J. M. Wyvratt et. al., *J. Org. Chem.*, **52**, 944-945 (1987)] followed by treatment with aqueous NaOH in DMF provides the alcohol **11**. Treatment of **11** with diazomethane followed by the reaction with CBr₄/Ph₃P yields the alkylating agent **12**.

A typical synthesis of alkylating agents bearing a substituted benzoxazole or benzthiazole ring is outlined in Scheme 5. The substituted benzoxazole **14** is prepared from the corresponding o-aminophenol **13** by the reaction of an appropriate orthoester under refluxing conditions (for other methods of synthesis of benzoxazoles see, S. A. Lang and Y. Lin, *Comprehensive Heterocyclic Chemistry*, Vol. 6, 1-130, Ed. C. W. Rees; and references cited therein). Reduction of **14** with NaBH₄ provides the alcohol **15** which is then subjected to pyridiniumdichromate oxidation (PDC) to yield the corresponding aldehyde **16**. Further elaboration of **16** as outlined provides the key

- 53 -


intermediate **17**. Similarly, the benzothiazole **19** can also be prepared from the appropriately substituted o-aminothiophenol **18**.

Scheme 4

- a. (I) DIBALH, Toluene; (II) Boc₂O, DMAP, CH₂Cl₂
b. (I) CHCl₃, KOH, DMF, 0°C; (II) NaOH, DME / H₂O
c. (I) CH₂N₂; (II) CBr₄/Ph₃P, CH₂Cl₂

- 54 -

Scheme 5



- 55 -

Scheme 6 illustrates the synthesis of benzofuran and dihydrobenzofuran alkylating agents **23** and **25**. The benzofuran **21** can be prepared from the α -phenoxy carbonyl compound **20** via a ring closure reaction [Stoermer and Wehln, Chem. Ber., **35**, 3549 (1902)] (for general methods of synthesis of benzofurans and dihydro-
5 benzofurans see, R. C. Elderfield and V. B. Meyer, Heterocyclic Compounds, Vol. 2, Chapter 1, Ed. R. C. Elderfield, Wiley; and references cited therein). The ester **21** is reduced to provide the aldehyde **22** which is then transformed into the corresponding
10 alkylating agent **23**. The dihydrobenzofuran ester **24**, obtained by catalytic reduction of **21**, can also be transformed into the corresponding alkylating agent **25** using the sequence of reactions outlined in Scheme 6.


Benzothiophene **26** may be synthesized from the
15 corresponding aldehyde **26b** in a manner similar to that outlined in Scheme 6 for benzofuran **23**. Benzothiophene **26b** can be prepared by the oxidative cyclization (using an alkaline solution of potassium ferricyanide) of appropriately substituted o-mercaptocinnamic acid **26a** [C. Chmielewsky and P. Friedlander, Chem. Ber., **46**, 1903
20 (1913)]. (For general methods of synthesis of benzothiophene, See, E. Champaigne in Comprehensive Heterocyclic Chemistry, vol. 4, Chapter 3-15; Eds. A. Katritzky and C.W. Rees.)

Scheme 7 outlines a typical synthesis of α -bromoarylacetates, **30** and **32**, bearing appropriately substituted
25 methylenedioxy or 1,4-dioxane rings. The substituted catechol derivative **27** is treated with an appropriate dibromide (where m is 1 or 2) in the presence of cesium carbonate in dimethylformamide to provide **28**. Treatment of **28** with DIBALH yields the aldehyde **29** which is then transformed into the desired alkyl bromide as described.

30


- 56 -

Scheme 6




- 57 -

Scheme 7

a. $\text{Br}-(\text{CH}_2)_m-\text{Br}$, Cs_2CO_3 , DMF

b. DIBALH, toluene

c. (i) CHCl_3 , KOH, DMF, 0°C ; (ii) NaOH, DME / H_2O ;
(iii) HCl / MeOH; (iv) $\text{CBr}_4/\text{Ph}_3\text{P}$, CH_2Cl_2 ;d. NBS, AIBN, CCl_4 

- 58 -

The reactions are performed in a solvent appropriate to the reagents and materials employed and suitable for the transformation being effected. It is understood by those skilled in the art of organic synthesis that the functionality present on the heterocycle and in the reactants being employed should be consistent with the chemical transformations being conducted. Depending upon the reactions and techniques employed, optimal yields may require changing the order of synthetic steps or use of protecting groups followed by deprotection.

The compounds useful in the novel method treatment of this invention form salts with various inorganic and organic acids and bases which are also within the scope of the invention. Such salts include ammonium salts, alkali metal salts like sodium and potassium salts, alkaline earth metal salts like the calcium and magnesium salts, salts with organic bases; e.g., dicyclohexylamine salts, N-methyl-D-glucamine, salts with amino acids like arginine, lysine, and the like. Also, salts with organic and inorganic acids may be prepared; e.g., HCl, HBr, H₂SO₄, H₃PO₄, methanesulfonic, toluenesulfonic, maleic, fumaric, camphorsulfonic.

The salts can be formed by conventional means, such as by reacting the free acid or free base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed *in vacuo* or by freeze-drying or by exchanging the cations of an existing salt for another cation on a suitable ion exchange resin.

It will be appreciated that the compounds of general Formula I in this invention may be derivatised at functional groups to provide prodrug derivatives which are capable of conversion back to the parent compounds *in vivo*. The concept of prodrug administration has been extensively reviewed (e.g. A.A. Sinkula in Annual Reports in Medicinal Chemistry, Vol 10, R.V. Heinzelman, Ed., Academic Press, New York London, 1975, Ch. 31, pp. 306-326), H. Ferres, Drugs of Today, Vol 19, 499-538 (1983) and J. Med. Chem., 18, 172 (1975). Examples of such prodrugs include the physiologically acceptable and metabolically labile ester derivatives, such as lower alkyl (e.g. methyl or ethyl esters), aryl

- 59 -

(e.g. 5-indanyl esters), alkenyl (e.g. vinyl esters), alkoxyalkyl (e.g. methoxymethyl esters), alkylthioalkyl (e.g. methylthiomethyl esters), alkanoyloxyalkyl (e.g. pivaloyloxymethyl esters), and substituted or unsubstituted aminoethyl esters (e.g. 2-dimethylaminoethyl esters).

5 Additionally, any physiologically acceptable equivalents of the compounds of general Formula I, similar to the metabolically labile esters, which are capable of producing the parent compounds of general Formula I *in vivo*, are within the scope of this invention.

It will be further appreciated that the majority of compounds
10 of general Formula I claimed herein are asymmetric and are produced as racemic mixtures of enantiomers and that both the racemic compounds and the resolved individual enantiomers are considered to be in the scope of this invention. The racemic compounds of this invention may be resolved to provide individual enantiomers utilizing methods known to
15 those skilled in the art of organic synthesis. For example, diastereoisomeric salts, esters or imides may be obtained from a racemic compound of general Formula I and a suitable optically active amine, amino acid, alcohol or the like. The diastereoisomeric salts, esters or imides are separated and purified, the optically active enantiomers are
20 regenerated and the preferred enantiomer is the more potent isomer. The resolved enantiomers of the compounds of general Formula I, their pharmaceutically acceptable salts and their prodrug forms are also included within the scope of this invention.

Endothelin (ET-1), and two closely related bioactive
25 peptides, ET-2 and ET-3, are widely distributed in mammalian tissues, and they can induce numerous biological responses in non-vascular as well as vascular tissues by binding to at least two distinct endothelin receptor subtypes. In addition to cardiovascular smooth muscle, neural and atrial sites, endothelin receptors may also be found in
30 gastrointestinal, kidney, lung, urogenital, uteral and placental tissues.

Endothelin is a potent vasoconstrictor peptide and thus plays an *in vivo* role in arterial pressure-volume homeostasis. Not only peripheral, but coronary vascular resistance as well, is increased by endothelin. Cardiac output is decreased, while plasma renin activity is

- 60 -

increased. There is a reduction in renal blood flow and glomerular filtration rate, while levels of atrial natriuretic factor, vasopressin, and aldosterone become elevated.

It is also considered, in accordance with the present invention, that antagonists for the endothelin receptor may be useful in preventing or reducing restenosis subsequent to denudation following angioplasty. Such denudation results from myointimal thickening following angioplasty, which is caused by increased endothelin release. Endothelin acts as a growth factor with respect to smooth muscle and fibroblastic cells, and possibly other types of cells, as well.

Endothelin is also a neuropeptide, acting on the posterior pituitary, where it modulates the release of the neurosecretory hormones vasopressin and oxytocin. Endothelin released from the posterior pituitary also acts as a circulating hormone, having a wide range of actions as discussed further above. This includes effects on the endocrine system, especially the adrenal glands. Endothelin increases plasma levels of epinephrine.

Consequently, the novel compounds of the present invention, which are receptor antagonists of endothelin, have therapeutic usefulness in preventing, decreasing or modulating the various physiological effects of endothelin discussed above, by wholly or partially blocking access of endothelin to its receptor.

Endothelin Receptor Binding Assays

The binding of the novel compound of this invention to the endothelin receptor was determined in accordance with the assay described in detail immediately below. It is similar to the assay described in Ambar *et al.* (1989) Biochem. Biophys. Res. Commun. 158, 195-201; and Kloog *et al.* (1989) FEBS Letters, 253, 199-202.

The endothelins (ETs) have a number of potent effects on a variety of cells, and exert their action by interacting with specific receptors present on cell membranes. The compounds described in the present invention act as antagonists of ET at the receptors. In order to

- 61 -

identify ET antagonists and determine their efficacy in vitro, the following three ligand receptor assays were established.

Receptor binding assay using cow aorta membrane preparation:

5 Thoracic aortae were obtained from freshly slaughtered calves and brought to the lab on wet ice. The adventitia were removed, and the aorta was opened up lengthwise. The luminal surface of the tissue was scrubbed with cheesecloth to remove the endothelial layer. The tissue was ground in a meat grinder, and suspended in ice-cold 0.25 M sucrose, 5 mM tris-HCl, pH 7.4, containing 0.5 µg/mL leupeptin and 7
10 µg/mL pepstatin A. Tissue was homogenized twice and then centrifuged for 10 minutes at 750 x g at 4°C. The supernatant was filtered through cheesecloth and centrifuged again for 30 minutes at 48,000 x g at 4°C. The membrane pellet thus obtained was resuspended in the buffer solution described above (including the protease inhibitors), and aliquots
15 were quick-frozen and stored at -70°C until use. Membranes were diluted into 50 mM KPi, 5 mM EDTA pH 7.5 containing 0.01 % human serum albumin. Assays were done in triplicate. Test compounds and 100 pM [¹²⁵I]-endothelin-1 (2000-2200 Ci/µmole, obtained from New England Nuclear or Amersham) were placed in a tube containing this
20 buffer, and the membranes prepared above were added last. The samples were incubated for 60 min at 37°C. At the end of this incubation, samples were filtered onto prewetted (with 2% BSA in water) glass fiber filter pads and washed with 150 mM NaCl, 0.1% BSA. The filters were
25 assayed for ¹²⁵I radioactivity in a gamma counter. Nondisplaceable binding of [¹²⁵I]-endothelin-1 was measured in the presence of 100 nM unlabelled endothelin-1 [Endothelin-1 (ET-1) was purchased from Peptides International (Louisville, KY)]. Specific binding is defined as total binding minus nondisplaceable binding. The inhibitory concentration (IC₅₀) which gives 50% displacement of the total
30 specifically bound [¹²⁵I]-endothelin-1 was presented as a measure of the potency of such compound as ET antagonist.

- 62 -

Receptor binding assay using rat hippocampal membrane preparation:

Rat hippocampi were obtained from freshly sacrificed male Sprague-Dawley rats and placed in ice cold 0.25 M sucrose, 5 mM tris-HCl, pH 7.4 containing 0.5 µg/mL leupeptin, 7 µg/mL pepstatin A.

5 Hippocampi were weighed and placed in a Dounce homogenizer with 25 volumes (wet weight to volume) ice-cold sucrose buffer in the presence of protease inhibitors. Hippocampi were homogenized using the Dounce (glass-glass) homogenizer with type A pestle, with the homogenizer immersed in ice. Tissue homogenate was centrifuged at 750 x g for 10 min at 4°C. Supernatant was filtered through dampened cheesecloth, and centrifuged again at 48,000 x g for 30 min at 4°C. Membrane pellets

10 were resuspended in sucrose buffer with protease inhibitors. Aliquots of this preparation were quick frozen and stored at -70°C until use. Membranes were diluted into 50 mM KPi, 5 mM EDTA pH 7.5 containing 0.01% human serum albumin. Assays were done in triplicate. Test compounds and 25 pM [¹²⁵I]-endothelin-1 (2000-2200 Ci/µmole, obtained from New England Nuclear or Amersham) were placed in a tube containing this buffer, and the membranes prepared above were added last. The samples were incubated for 60 min at 37°C. At the end of this incubation, samples were filtered onto prewetted (with 2% BSA in water)

20 glass fiber filter pads and washed with 150 mM NaCl, 0.1% BSA. The filters were assayed for ¹²⁵I radioactivity in a gamma counter. Nondisplaceable binding of [¹²⁵I]-endothelin-1 was measured in the presence of 100 nM unlabelled endothelin-1 [Endothelin-1 (ET-1) was purchased from Peptides International (Louisville, KY)]. Specific binding is defined as total binding minus nondisplaceable binding. The inhibitory concentration (IC₅₀) which gives 50% displacement of the total specifically bound [¹²⁵I]-endothelin-1 was presented as a measure of the potency of such compounds as endothelin antagonists.

30 Receptor binding assay using cloned human ET receptors expressed in Chinese Hamster Ovary Cells:

Both endothelin receptor subtypes were cloned from a human cDNA library and were individually expressed in Chinese

- 63 -

Hamster Ovary cells. Cells were harvested by addition of 126 mM NaCl, 5 mM KCl, 2 mM EDTA, 1 mM NaH₂PO₄, 15 mM glucose, 10 mM tris/HEPES pH 7.4. Cells were centrifuged at 250 x g for 5 minutes. The supernatant was aspirated off, and the cells were resuspended in the 50 mM KPi, 5 mM EDTA pH 7.5 containing 0.01% human serum albumin. Assays were done in triplicate. Test compounds and 25-100 pM [¹²⁵I]-endothelin-1 (2000-2200 Ci/μmole, obtained from New England Nuclear or Amersham) were placed in a tube containing 50 mM KPi, 5 mM EDTA pH 7.5 containing 0.01% human serum albumin, and the cells prepared above were added last. The samples were incubated for 60 min at 37°C. At the end of this incubation, samples were filtered onto prewetted (with 2% BSA in water) glass fiber filter pad and washed with 150 mM NaCl, 0.1% BSA.

The filters were assayed for ¹²⁵I radioactivity in a gamma counter. Nondisplaceable binding of [¹²⁵I]-endothelin-1 was measured in the presence of 100 nM unlabelled endothelin-1 [Endothelin-1 (ET-1) was purchased from Peptides International (Louisville, KY)]. Specific binding is defined as total binding minus nondisplaceable binding. The inhibitory concentration (IC₅₀) which gives 50% displacement of the total specifically bound [¹²⁵I]-endothelin-1 was presented as a measure of the potency of such compounds as endothelin antagonists.

The binding assays described above were used to evaluate the potency of interaction of the compounds of the invention with endothelin receptors. To determine whether these compounds were endothelin antagonists, assays which measure the ability of the compounds to inhibit endothelin-stimulated phosphatidylinositol hydrolysis were established. Rat uterus contains predominantly one of the known endothelin receptor subtypes (ET_A).

Phosphatidylinositol hydrolysis assays using rat uterine slices:

Diethylstilbestrol primed female Sprague-Dawley rats were sacrificed and their uteri were collected, dissected of fat and connective tissue and minced. Minced tissue was added to oxygenated (95% O₂, 5% CO₂) 127 mM NaCl, 25 mM NaHCO₃, 10 mM Glucose, 2.5 mM KCl,

- 64 -

1.2 mM KH_2PO_4 , 1.2 mM MgSO_4 , 1.8 mM CaCl_2 . To the tissue mince, 1.2 μM myo-[^3H]-inositol (Amersham) was added. The mince was incubated 90 min at 37°C , with constant oxygenation. After incubation, the loaded tissue mince was washed five times with the same oxygenated buffer to remove excess radiolabelled inositol. The tissue mince was resuspended in the above buffer, containing 10 mM LiCl , aliquotted into tubes, and 3 nM endothelin-1 with and without test compounds was added to a final concentration of 3 nM to start the assay. Assays were done in quadruplicate. Samples were incubated at 37°C under blowing O_2 in a hooded water bath for 30 minutes. Reaction was stopped by addition of trichloroacetic acid to 6% concentration. Samples were sonicated for 10 min, centrifuged 20 min, then trichloroacetic acid was extracted with water-saturated ethyl ether. An aliquot of each sample was neutralized and diluted by addition of 50 mM tris-HCl pH 7.4. A 100 μL aliquot of this solution was assayed for radioactivity in a beta counter. The diluted neutralized sample was applied to Dowex 1 x 8-formate columns, washed with water, then washed with 60 mM ammonium formate, 5 mM sodium tetraborate. Samples were eluted with 200 mM ammonium formate, 5 mM sodium tetraborate. The radioactivity of each eluted sample was measured in a beta counter. Radioactivity was normalized by dividing radioactivity in post column sample by radioactivity in precolumn sample. Control values (100% stimulated) are values in the presence of endothelin minus the values in the absence of endothelin (basal). Test sample values are the values in the presence of endothelin and test sample minus basal. Inhibitory concentration (IC_{50}) is the concentration of test compound required to give a sample activity of 50% of control value.

Sarafotoxin S6c is a member of the endothelin family which binds preferentially to one of the known endothelin receptor subtypes (ET_B).

Phosphatidylinositol hydrolysis assays using rat lung slices:

Male Sprague-Dawley rats were sacrificed and their lungs were collected, dissected of fat and connective tissue and minced. Minced tissue was added to oxygenated (95% O_2 , 5% CO_2) 127 mM

- 65 -

NaCl, 25 mM NaHCO₃, 10 mM Glucose, 2.5 mM KCl, 1.2 mM KH₂PO₄, 1.2 mM MgSO₄, 1.8 mM CaCl₂. To the tissue mince, 1.2 μM myo-[³H]-inositol was added. The mince was incubated 60 min at 37°C, with constant oxygenation. After incubation, loaded tissue mince was
5 washed five times with the same oxygenated buffer to remove excess radiolabelled inositol. Tissue mince was resuspended in the above buffer, containing 10 mM LiCl, aliquotted into tubes, and sarafotoxin S6c (to a final concentration of 3 nM) with and without test compounds was added to start the assay. Assays were done in quadruplicate. Samples were
10 incubated at 37°C under blowing O₂ in a hooded water bath for 30 minutes. Reaction was stopped by addition of 0.5 mL 18% trichloroacetic acid to 6% concentration. Samples were sonicated for 10 min, centrifuged 20 min, then trichloroacetic acid was extracted with water-saturated ethyl ether. An aliquot of each sample was neutralized
15 and diluted by addition of 50 mM tris-HCl pH 7.4. A 100 μL aliquot of this solution was assayed for radioactivity in a beta counter. The diluted neutralized sample was applied to Dowex 1 x 8-formate columns, washed with water, then washed with 60 mM ammonium formate, 5 mM sodium tetraborate. Samples were eluted with 200 mM ammonium formate, 5
20 mM sodium tetraborate. The radioactivity of each eluted sample was measured in a beta counter. Radioactivity was normalized by dividing radioactivity in post column sample by radioactivity in precolumn sample. Control values (100% stimulated) are values in the presence of sarafotoxin minus the values in the absence of sarafotoxin (basal). Test
25 sample values are the values in the presence of sarafotoxin and test sample minus basal. Inhibitory concentration (IC₅₀) is the concentration of test compound required to give a sample activity of 50% of control value.

30 Phosphatidylinositol hydrolysis assays using cloned human endothelin receptors expressed in Chinese Hamster Ovary cells:

Endothelin receptors of both receptor subtypes were cloned from a human cDNA library and were individually expressed in Chinese Hamster Ovary cells. Cells were loaded overnight by the addition of 1.2

- 66 -

5 μ M myo-[3 H]-inositol to their growth medium. Cells were harvested by addition of 126 mM NaCl, 5 mM KCl, 2 mM EDTA, 1 mM NaH_2PO_4 , 15 mM glucose, 10 mM tris/HEPES pH 7.4. Cells were washed five times by centrifugation at 250 x g for 5 minutes to remove excess radiolabelled inositol. The supernatant was aspirated off, and the cells were resuspended in the same oxygenated (95% O_2 , 5% CO_2) buffer containing 10 mM LiCl, aliquotted into tubes, and endothelin-1 (to a final concentration of 0.3 nM) with and without test compounds was added to start the assay. Assays were done in quadruplicate. Samples were incubated at 37°C under blowing O_2 in a hooded water bath for 30 minutes. Reaction was stopped by addition of 0.5 mL 18% trichloroacetic acid to 6% concentration. Samples were sonicated for 10 min, centrifuged 20 min, then trichloroacetic acid was extracted with water-saturated ethyl ether. An aliquot of each sample was neutralized and diluted by addition of 50 mM tris-HCl pH 7.4. A 100 μ L aliquot of this solution was assayed for radioactivity in a beta counter. The diluted neutralized sample was applied to Dowex 1 x 8-formate columns, washed with water, then washed with 60 mM ammonium formate, 5 mM sodium tetraborate. Samples were eluted with 200 mM ammonium formate, 5 mM sodium tetraborate. The radioactivity of each eluted sample was measured in a beta counter. Radioactivity was normalized by dividing radioactivity in post column sample by radioactivity in precolumn sample. Control values (100% stimulated) are values in the presence of endothelin minus the values in the absence of endothelin (basal). Test sample values are the values in the presence of endothelin and test sample minus basal. Inhibitory concentration (IC_{50}) is the concentration of test compound required to give a sample activity of 50% of control value.

Using the methodology described above, the compounds of the invention were evaluated and found to exhibit IC_{50} values of at least <50 μ M thereby demonstrating and confirming the utility of the compounds of this invention as an effective endothelin antagonists.

Accordingly the novel compounds of the present invention are useful in human therapy for treating asthma, hypertension, pulmonary hypertension, arteriosclerosis, heart failure, renal failure particularly

- 67 -

post-ischemic renal failure, cyclosporin nephrotoxicity, vasospasm, vascular restenosis, cerebral and cardiac ischemia and other ischemic states, myocardial infarction, Raynaud's disease, inflammatory bowel diseases, including Crohn's disease and ulcerative colitis, as well as other
5 inflammatory diseases, or endotoxic shock caused by or associated with endothelin, by administration to a patient in need of such treatment of a therapeutically effective amount thereof.

In the management of hypertension and the clinical conditions noted above, the compounds of this invention may be utilized
10 in compositions such as tablets, capsules or elixirs for oral administration, suppositories for rectal administration, sterile solutions or suspensions for parenteral or intramuscular administration, and the like. The compounds of this invention can be administered to patients (animals and human) in need of such treatment in dosages that will provide optimal
15 pharmaceutical efficacy. Although the dose will vary from patient to patient depending upon the nature and severity of disease, the patient's weight, special diets then being followed by a patient, concurrent medication, and other factors which those skilled in the art will recognize, the dosage range will generally be about 0.5 mg.- 1.0 g. per
20 patient per day which can be administered in single or multiple doses. Preferably, the dosage range will be about 0.5 - 500 mg. per patient per day; more preferably about 0.5 - 100 mg. per patient per day.

The present invention also relates to pharmaceutical compositions for treating asthma, hypertension, pulmonary hypertension, arteriosclerosis, heart failure, renal failure particularly post-ischemic
25 renal failure, cyclosporin nephrotoxicity, vasospasm, vascular restenosis, cerebral and cardiac ischemia and other ischemic states, myocardial infarction, Raynaud's disease, inflammatory bowel diseases, including Crohn's disease and ulcerative colitis, as well as other inflammatory
30 diseases, or endotoxic shock caused by or associated with endothelin, comprising a therapeutically effective amount of the novel compound of this invention together with a pharmaceutically acceptable carrier thereof.

About 0.5 mg.- 1.0 g. of compound or mixture of compounds of Formula I or a physiologically acceptable salt is

- 68 -

compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in a unit dosage form as called for by accepted pharmaceutical practice. The amount of active substance in these compositions or preparations is such that a suitable dosage in the range indicated is obtained.

5 Illustrative of the adjuvants which can be incorporated in tablets, capsules and the like are the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; an excipient such as microcrystalline cellulose; a disintegrating agent such as corn starch, pregelatinized starch, alginic acid and the like; a lubricant such as 10 magnesium stearate; a sweetening agent such as sucrose, lactose or saccharin; a flavoring agent such as peppermint, oil of wintergreen or cherry. When the dosage unitform is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as fatty oil. Various other materials may be present as coatings or to otherwise 15 modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propyl parabens as preservatives, a dye and a flavoring such as cherry or orange 20 flavor.

Sterile compositions for injection can be formulated according to conventional pharmaceutical practice by dissolving or suspending the active substance in a vehicle such as water for injection, a naturally occurring vegetable oil like sesame oil, coconut oil, peanut oil, cottonseed oil, etc., or a synthetic fatty vehicle like ethyl oleate or the 25 like. Buffers, preservatives, antioxidants and the like can be incorporated as required.

The following examples illustrate the preparation of the compounds of Formula I and their incorporation into pharmaceutical 30 compositions and as such are not to be considered as limiting the invention set forth in the claims appended hereto.

- 69 -

EXAMPLE 1

2-Butyl-3-[4-((1-carboxy-1-phenyl)methoxy)-3-allyl]phenyl]methyl-6-methylquinazolin-4(3H)-one

5

Step A: Preparation of 4-(2-propen-1-yloxy)benzyl alcohol

10

To a suspension of NaH (130 mg; 4.33 mmol) in DMF (5 mL) at 0°C under nitrogen was added a solution of 4-hydroxymethylphenol (512 mg; 4.12 mmol) in DMF (5 mL). After stirring 5 minutes at room temperature, a solution of allyl bromide (375 mL, 4.33 mmol) in DMF (5 mL) was added dropwise. The reaction was stirred for 20 minutes at 0°C, then quenched with water and concentrated *in vacuo*.

15

The residue was partitioned between water and ethyl acetate. The combined organic layers were washed with 4% HCl, saturated NaHCO₃, and then brine, and dried (MgSO₄), filtered and concentrated *in vacuo* to yield 650 mg (97%) of the title compound.

20

¹H NMR (300 MHz, CDCl₃, ppm): δ 7.27 (dd, 2H), 6.90 (apparent d, 2H), 6.12-5.98 (m, 1H), 5.41 (apparent dd, 1H), 5.29 (dd, 1H), 4.58 (s, 2H), 4.52 (dd, 2H), 1.93 (br s, 1H).

25

Step B: Preparation of 4-tert-butyldimethylsilyloxymethylphenyl-(2-propen-1-yl) ether

30

To a solution of the product of Step A (650 mg, 3.96 mmol) in CH₂Cl₂ (20 mL), cooled to 0°C, under nitrogen, was added triethylamine (612 mL, 4.39 mmol) and a solution of tert-butyldimethylsilyl chloride (631 mg; 4.19 mmol) in CH₂Cl₂ (2 mL). After stirring at room temperature for 18 hours, the reaction mixture was diluted with ethyl acetate (60 mL), washed with water, and saturated sodium bicarbonate, and then dried (MgSO₄). The filtrate was concentrated *in vacuo* to afford 1.1 g of the title compound which was used crude in the next reaction (R_f=0.45, 5% ethyl acetate/hexanes).

- 70 -

^1H NMR (300 MHz, CDCl_3 , ppm): δ 7.24 (d, 2H), 6.89 (d, 2H), 6.15-6.00 (m, 1H), 5.42 (apparent d, 1H), 5.30 (apparent d, 1H), 4.68 (s, 2H), 4.53 (apparent dd, 2H), 0.95 (s, 9H), 0.12 (s, 6H).

5 Step C: Preparation of 4-tert-butyl dimethylsilyloxymethyl-2-allyl-phenol

The product of Step B (0.51 g, 1.83 mmol) was heated to 200°C under a nitrogen atmosphere for 5 hours. The crude reaction mixture was dissolved in eluant and chromatographed on silica (MPLC, 10 5/95 ethyl acetate/hexane) to afford 178 mg (35%) of the title compound (R_f = 0.11, 5% ethyl acetate/hexanes).

^1H NMR (300 MHz, CDCl_3 , ppm): δ 7.10 (unsymmetrical d, 2H), 6.78 (d, 2H), 6.09-5.94 (m, 2H), 5.21-5.11 (m, 2H), 5.00 (s, 1H), 4.67 (s, 2H), 15 3.40 (d, 2H), 0.95 (s, 9H), 0.11 (s, 6H).

FAB MS: m/e = 277 ($M+1$).

Step D: Preparation of methyl 2-(4-tert-butyl-dimethylsilyloxy-methyl-2-allylphenoxy)-2-phenylacetate

20 To a suspension of KH (1.3 eq) in DMF (1 mL) was added a solution of the product of Step C (157 mg, 0.566 mmol) in DMF (1 mL), followed by 18-crown-6 (30 mg; 0.2 eq). The reaction mixture was stirred for 5 minutes at room temperature. A solution of methyl 2-bromophenylacetate (168 mg, 0.735 mmol) in DMF (1 mL) was added, 25 followed by a catalytic amount of potassium iodide. The reaction was heated to 80°C for 0.5 hours then stirred at room temperature for 16 hours. After concentration *in vacuo*, the residue was partitioned between water and ethyl acetate. The combined organic layers were washed with water, brine, then dried (MgSO_4). After filtration and concentration *in* 30 *vacuo*, the residue was chromatographed on silica (MPLC, ethyl acetate/hexanes (5/95)) to afford 158 mg (66%) of the title compound (R_f =0.22, 5% ethyl acetate/hexanes).

- 71 -

¹H NMR (300 MHz, CDCl₃, ppm): δ 7.64-7.57 (dd, 2H), 7.46-7.35 (m, 3H), 7.17-7.06 (m, 2H), 6.72 (d, 1H), 6.12-5.98 (m, 1H), 5.65 (s, 1H), 5.11-5.04 (m, 2H), 4.66 (s, 2H), 3.72 (s, 3H), 3.53 (d, 2H), 0.95 (s, 9H), 0.10 (s, 6H).

5

FAB MS: consistent with structure.

Step E: Preparation of methyl 2-(4-bromomethyl-2-allylphenoxy)-2-phenylacetate

10

To a cooled (0°C) solution of the product of Step D (156 mg, 0.366 mmol) in CH₃CN (2 mL), were added carbon tetrabromide (182 mg, 0.55 mmol) and triphenylphosphine (144 mg, 0.55 mmol). After 30 minutes at 0°C, the reaction mixture was allowed to warm to room temperature, at which point acetone (40 mL, 0.55 mmol) was added. After 16 hours at room temperature, the reaction mixture was filtered, the filtrate was concentrated *in vacuo*, and the residue was purified on a silica gel flash chromatography column eluted with 5% ethyl acetate/hexanes to afford 86 mg (63 %) of the title compound (R_f=0.13, 5% ethyl acetate/hexanes).

15

20

¹H NMR (300 MHz, CDCl₃, ppm): δ 7.67-7.57 (dd, 2H), 7.47-7.37 (m, 3H), 7.27-7.13 (m, 2H), 6.72 (d, 1H), 6.16-5.98 (m, 1H), 5.68 (s, 1H), 5.20-5.08 (m, 2H), 4.49 (s, 2H), 3.73 (s, 3H), 3.54 (d, 2H).

FAB MS: consistent with structure.

25

Step F: Preparation of 2-n-butyl-6-methylquinazolin-4(1H)-one

30

To a solution of 3.0 g (20 mmol) of 2-amino-5-methyl benzoic acid in 20 mL of dry DMF at 0°C was added 200 mg of DMAP followed by 6.07 g (60 mmol) of triethylamine and 5.02 g (40 mmol) of valeryl chloride. The resulting mixture was stirred at 0°C for 30 min. The mixture was heated to 110°C and monitored by TLC for the formation of the intermediate quinoxazolone (R_f=0.8, 40% EtOAc/hexane). Following complete formation of the intermediate 10 g (100 mmol) of (NH₄)₂CO₃ was added cautiously. Heating was continued to

- 72 -

ensure consumption of the quinoxazolone and formation of the polar ($R_f=0.4$, 40% EtOAc/hexane) quinazolin-4(1H)-one. The reaction mixture was concentrated *in vacuo* and the residue was taken up in 50 mL of ether and 50 mL of water. The mixture was filtered and the filtrate discarded after washing the residue with 20 mL of ether. The residue was recrystallized from MeOH to give 1.07 g (25%) of the title compound as a white crystalline solid.

^1H NMR (300 MHz, CDCl_3 , ppm): δ 0.94 (t, 3H, $J=6.7$ Hz), 1.50 (m, 2H), 1.83 (m, 2H), 2.49 (s, 3H), 2.78 (t, 2H), 7.60 (m, 2H), 8.05 (m, 1H).

Anal ($\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}$), C, H, N.

Step G: Preparation of 2-Butyl-3-[4-((1-carbomethoxy-1-phenyl)-methoxy)-3-allylphenyl]methyl-6-methylquinazolin-4(3H)-one

To a suspension of NaH (0.514 mmol) in DMF (2 mL) was added 92 mg (0.428 mmol) of 2-butyl-6-methylquinazolin-4(3H)-one (Step F of Example 1) and the reaction mixture was stirred for 30 minutes at room temperature. A solution of the product of Step E (177 mg, 0.471 mmol) in DMF (1.5 mL) was added, and the reaction was stirred at room temperature for 18 hours. The reaction mixture was then concentrated *in vacuo* and partitioned between water and ethyl acetate. The combined organic layers were washed with water then brine, dried (MgSO_4), filtered, and concentrated *in vacuo*. The residue was chromatographed on silica (MPLC, hexanes/ethyl acetate (4/1)) to afford 96 mg (44 %) of the title compound.

^1H NMR (300 MHz, CDCl_3 , ppm): δ 8.08 (s, 1H), 7.6-7.49 (br s, 4H), 7.42-7.28 (m, 3H), 7.09-7.01 (br s, 1H), 6.93-6.83 (br dd, 1H), 6.66 (d, 1H), 6.08-5.92 (m, 1H), 5.60 (s, 1H), 5.36-5.22 (br s, 2H), 5.12-4.98 (m, 2H), 3.68 (s, 3H), 3.48 (d, 2H), 3.48 (d, 2H), 2.72 (t, 2H), 2.48 (s, 3H), 1.80-1.65 (m, 2H), 1.40 (q, 2H), 0.90 (t, 3H).

FAB MS: $m/e = 511$ ($M+1$).

- 73 -

Step H: Preparation of 2-Butyl-3-[4-((1-carboxy-1-phenyl)-methoxy)-3-allylphenyl]methyl-6-methylquinazolin-4(3H)-one

5 To a solution of the product of Step G (20 mg, 0.039 mmol) in MeOH (2 mL), were added 4 drops of water and 2.0 N NaOH (22 mL, 0.043 mmol). After stirring for 18 hours at room temperature, the reaction mixture was concentrated *in vacuo*, dissolved in water/THF, and treated with HCl (0.15 mL, 1.0 N) at room temperature for 30 minutes. The reaction mixture was then concentrated *in vacuo* and chromatographed on a Sephadex LH-20 column eluted with MeOH to afford 19 mg of crude product, which was recrystallized from MeOH to yield 2 mg (10%) of the title compound.

15 ¹H NMR (300 MHz, CD₃OD, ppm): δ 8.05 (s, 1H), 7.70 (d, 1H), 7.63-7.48 (m, 3H), 7.44-7.28 (m, 3H), 7.08 (s, 1H), 6.99-6.92 (m, 1H), 6.83 (d, 1H), 6.07-5.92 (m, 3H), 5.73 (s, 1H), 5.40 (s, 2H), 5.08-5.88 (m, 2H), 3.46 (br s, 2H), 2.80 (t, 2H), 2.51 (s, 3H), 1.68-1.57 (m, 2H), 1.43-1.28 (m, 2H), 0.88 (t, 3H).

20 FAB MS: m/e = 497 (M+1).

EXAMPLE 2

25 2-Butyl-3-[4-((1-carboxy-1-phenyl)methoxy)-3-propylphenyl]methyl-6-methylquinazolin-4(3H)-one

Step A: Preparation of 2-Butyl-3-[4-((1-carbomethoxy-1-phenyl)-methoxy)-3-propylphenyl]methyl-6-methylquinazolin-4(3H)-one

30 To a solution of the product of Example 1, Step G (20 mg, 0.039 mmol) in CH₂Cl₂ (2 mL), was added Wilkinson's catalyst (7.6 mg). The reaction mixture was hydrogenated at 40 psi, room temperature for 4.5 hours. After concentration *in vacuo*, the residue was

- 74 -

chromatographed on silica (MPLC, hexanes/ethyl acetate (4/1) to afford 15 mg (78%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 8.10 (s, 1H), 7.63-7.54 (m, 4H), 7.46-7.36 (m, 3H), 7.02 (apparent s, 1H), 6.89 (dd, 1H), 6.63 (d, 1H), 5.62 (s, 1H), 5.32 (br s, 2H), 3.71 (s, 3H), 2.79-2.63 (m, 4H), 2.50 (s, 3H), 1.80-1.60 (m, 2H), 1.48-1.34 (m, 2H), 1.02-0.87 (m, 6H).

FAB-MS: m/e = 513 (M+1).

10 Step B: Preparation of 2-Butyl-3-[4-((1-carboxy-1-phenyl)-methoxy)-3-propylphenyl]methyl-6-methylquinazolin-4(3H)-one

To a solution the product of Step A (11 mg; 0.22 mmol) in MeOH (2 mL) was added NaOH (1.5 eq, 2.0 N) and a few drops of water. After stirring for 20 hours at room temperature, the reaction mixture was concentrated *in vacuo*, dissolved in water/THF, treated with HCl (5 eq) for 30 minutes at room temperature, concentrated *in vacuo*, and chromatographed on a Sephadex LH-20 column eluting with MeOH to afford 11 mg (99%) of the title compound.

20 ¹H NMR (300 MHz, CD₃OD, ppm): δ 8.19 (s, 1H), 7.88 (d, 1H), 7.68 (d, 1H), 7.58 (dd, 2H), 7.48-7.33 (m, 3H), 7.18 (s, 1H), 7.08 (d, 1H), 6.84 (d, 1H), 5.70 (s, 1H), 5.48 (s, 2H), 3.15-3.03 (m, 2H), 2.69 (t, 2H), 2.53 (s, 3H), 1.72-1.5 (m, 4H), 1.49-1.34 (m, 2H), 0.98-0.84 (m, 6H).

25 FAB-MS: m/e = 499 (M+1).

EXAMPLE 3

30 2-Butyl-3-[4-((1-carboxy-1-phenyl)methoxy)-3-chloro-phenyl]methyl-6-methylquinazolin-4(3H)-one

Step A: Preparation of methyl 2-(2-chloro-4-methyl-phenoxy)-2-phenylacetate

- 75 -

To a suspension of 0.282 g (7.04 mmol) of a 60% oil dispersion of sodium hydride in DMF was added 1.00 g (7.04 mmol) of 2-chloro-4-methylphenol and the mixture was stirred under an N₂ atmosphere at room temperature. After 10 minutes, a solution of 1.94 g (8.45 mmol) of methyl 2-bromophenylacetate dissolved in 10 mL of DMF was added and the reaction was stirred an additional 1.5 hours. The reaction was then diluted into ethyl acetate, washed with water, dried (MgSO₄), filtered and evaporated. The residue was purified on a silica gel flash chromatography column eluted with 4% ethyl acetate/hexane to afford 1.70 g (83%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 2.20 (s, 3H), 3.70 (s, 3H), 5.60 (s, 1H), 6.70-6.80 (d, 1H), 6.85-6.95 (d, 1H), 7.20 (br s, 1H), 7.20-7.30 (m, 3H), 7.55-7.65 (m, 2H).

EI-MS: m/e 290 (M⁺).

Step B: Preparation of methyl 2-(2-chloro-4-bromo-methyl-phenoxy)-2-phenylacetate

To a solution of 1.70 g (5.86 mmol) of the product from Step A dissolved in 20 mL of CCl₄ was added 1.04 g (5.86 mmol) of N-bromosuccinimide and 50 mg (catalytic amount) of AIBN. The reaction mixture was stirred and heated at reflux for 7 hours, then an additional 0.20 g of NBS was added. The reaction was refluxed for 48 hours, then cooled and concentrated *in vacuo*. The residue was purified on a silica gel flash chromatography column eluted with 10% ethyl acetate/hexane to afford 0.730 g (34%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 3.70 (s, 3H), 4.40 (s, 2H), 5.65 (s, 1H), 6.75-6.85 (d, 1H), 7.10-7.20 (d, 1H), 7.30-7.45 (m, 4H), 7.55-7.65 (m, 2H).

FAB-MS: m/e 369 (M+1).

- 76 -

Step C: Preparation of 2-butyl-3-[4-((1-carbomethoxy-1-phenyl-methoxy)-3-chlorophenyl)methyl-6-methylquinazolin-4(3H)-one]

To a half suspension of 62 mg (0.287 mmol) of the product of Step F of Example 1 in 1.0 mL of anhydrous DMF was added 12 mg (1.05 eq) of a 60% oil dispersion of sodium hydride and the reaction mixture was stirred under an N₂ atmosphere. After stirring 10 minutes at room temperature, a solution of 0.127 g (1.2 eq) of the product of Step B dissolved in 1.0 mL DMF was added to the solution of the anion. The reaction mixture was then stirred overnight, then partitioned between water and ethyl acetate. The organic layer was separated, dried (MgSO₄), filtered and evaporated *in vacuo*. The residual oil was purified on a silica gel flash chromatography column eluted with 30% ethyl acetate/hexane to afford 76 mg (52%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 0.87-0.94 (t, 3H), 1.32-1.45 (m, 2H), 1.67-1.78 (m, 2H), 2.42 (s, 3H), 2.66-2.72 (t, 2H), 3.70 (s, 3H), 5.30 (br s, 2H), 5.60 (s, 1H), 6.77 (d, 1H), 6.94 (dd, 1H), 7.18 (s, 1H), 7.33-7.42 (m, 3H), 7.53-7.61 (m, 4H), 8.06 (s, 1H).

FAB-MS: m/e 505, 507 (M+1, 3:1 ratio).

Step D: Preparation of 2-butyl-3-[4-((1-carboxy-1-phenyl)methoxy)-3-chlorophenyl)methyl-6-methylquinazolin-4(3H)-one]

To a solution of 72 mg of the product of Step C dissolved in 2 mL of methanol, was added 0.25 mL of a 1 N solution of NaOH and the reaction mixture was stirred at room temperature for 3 days. The reaction mixture was then adjusted to pH 7 with 1 N HCl, concentrated *in vacuo*, and the residue was applied to a silica gel flash chromatography column and eluted with CHCl₃/MeOH/NH₄OH (80:15:1). Evaporation of the purified fractions and drying *in vacuo* afforded 50 mg (71%) of the title compound.

¹H NMR (300 MHz, CD₃OD, ppm): δ 0.92-0.98 (t, 3H), 1.37-1.48 (m, 2H), 1.66-1.77 (m, 2H), 2.54 (s, 3H), 2.78-2.84 (t, 2H), 5.40 (s, 2H), 5.68

- 77 -

(s, 1H), 7.01-7.10 (m, 2H), 7.31-7.44 (m, 4H), 7.58-7.72 (m, 4H), 8.07 (s, 1H).

FAB-MS: m/e 491, 493 (M+1, 3:1 ratio).

5

EXAMPLE 4

3-[4-((1-Carboxy-1-phenyl)methoxy)-3-chloro-5-methoxyphenyl]methyl-6-(N-methyl-N-iso-butyloxycarbonyl)amino-2-propylquinazolin-4(3H)-one

10

Step A: Preparation of 2-propyl-6-nitroquinazolin-4(1H)-one

15

To a suspension of 48.94 g (0.3 mol) of 3-nitro-5-amino-benzonitrile in 500 mL of CH₂Cl₂ was added 63 mL of Et₃N, 3 g DMAP and lastly, dropwise, 45.5 g (0.45 mol) of butyryl chloride. A mild exothermic reaction ensued. The mixture was allowed to stir for 2 days (monitored by TLC with 50% EtOAc/hexanes). The solution was washed with 1 N HCl (2 x 100 mL), water (1 x 100 mL), saturated NaHCO₃ (2 x 100 mL), brine (1 x 100 mL) and dried over MgSO₄. The suspension was filtered and concentrated *in vacuo*. The residue was suspended in a mixture of 600 mL of MeOH and 200 mL of water in a three neck round bottom flask. To this was added gradually 140 mL of 5 N NaOH (0.7 mol) solution followed by the dropwise addition of 80 mL of 30% H₂O₂ (0.7 mmol) solution (exothermic). The mixture was refluxed overnight, cooled to room temperature and filtered. The filtrate was acidified with 1 N HCl cooled to 5°C and filtered. The quinazolinone was recrystallized from hot MeOH to give 38 g (54%) of the title compound as pale brown fine crystals.

20

25

¹H NMR (300 MHz, CDCl₃, ppm): δ 1.10 (t, 3H, J=7.8 Hz), 1.93 (m, 2H), 2.79 (t, 2H, J=7.3 Hz), 7.81 (d, 1H, J=8.9 Hz), 8.55 (dd, 1H, J=2.4, 8.8 Hz), 9.14 (d, 1H, J=2.4 Hz), 10.72 (br s, 1H).

30

Step B: Preparation of 3-(4,4'-dimethoxybenzhydryl)-2-propyl-6-nitroquinazolin-4(3H)-one

- 78 -

To a suspension of 1.01 g (33.7 mmol) of 80% sodium hydride in 20 mL of dry DMF was added at 0°C 7.5 g (32 mmol) of the product of Step A as a solid. The reaction mixture was diluted with a further 50 mL of DMF to assist stirring. After hydrogen evolution was complete, a solution of 8.8 g (33.7 mmol) of 4,4'-dimethoxybenzhydryl chloride in 20 mL of dry DMF was added dropwise. The reaction mixture was stirred overnight and then poured into 300 mL of 0.1 N NaOH. The precipitate was collected by filtration and dried under vacuum to give 12.1 g (94%) of a yellow solid.

¹H-NMR (300 MHz, CDCl₃, ppm): δ 0.87 (t, 3H, J=7.3 Hz), 1.58 (br m, 2H), 2.72 (t, 2H, J=7.8 Hz), 3.80 (s, 6H), 6.88 (d, 4H, J=9 Hz), 7.19 (d, 4H, J=9.0 Hz), 7.73 (d, 1H, J=8.9 Hz), 8.48 (dd, 1H, J=2.8, 9.0 Hz), 9.08 (d, 1H, J=2.8 Hz).

Step C: Preparation 6-amino-3-(4,4'-dimethoxybenz-hydryl)-2-propylquinazolin-4(3H)-one

A solution of 12.1 g (26.0 mmol) of the product of Step B dissolved in 250 mL of EtOAc was hydrogenated under atmospheric pressure over three days in the presence of three portions of 1.2 g of 10% Pd/C added daily. The mixture was filtered through celite and concentrated *in vacuo* to give an oil. The product was purified by flash chromatography over silica gel eluted with 50% EtOAc/hexanes to give 7.8 g (72%) of the amine.

¹H-NMR (300 MHz, CDCl₃, ppm): δ 0.82 (t, 3H, J=7.2 Hz), 1.49 (br m, 2H), 2.61 (t, 2H, J=7.81 Hz), 3.79 (s, 6H), 3.90 (br s, 2H), 6.85 (d, 4H, J=8.8 Hz), 7.08 (dd, 1H, J=2.8, 8.7 Hz), 7.20 (d, 4H, J=8.4 Hz), 7.42 (d, 1H, J=2.7 Hz), 7.47 (d, 1H, J=8.7 Hz).

Step D: Preparation of 3-(4,4'-dimethoxybenzhydryl)-6-(N-methyl-N-isobutyloxycarbonyl)-amino-2-propyl-quinazolin-4(3H)-one

To a suspension of 81.5 mg (2.7 mmol) of 80% NaH in 3 mL of dry DMF at 0°C under nitrogen was added dropwise a solution of

- 79 -

1.03 g (2.5 mmol) of 6-amino-3-(4,4'-dimethoxybenzhydryl)-2-propylquinazolin-4(3H)-one dissolved in 3 mL of DMF. The resulting mixture was stirred for 30 minutes and then treated with 0.35 mL (2.7 mmol) of neat isobutylchloroformate. The solution was stirred for 30 minutes and then treated with 2.97 mL (2.97 mmol) of a 1 M solution of lithium bis(trimethylsilyl)amide in THF. The dark solution was stirred for a further 30 minutes at 0°C and then was treated with 0.2 mL (3.26 mmol) of neat iodomethane. The mixture was stirred overnight at room temperature, poured into 50 mL of EtOAc and washed consecutively with water (2x10 mL), brine (1x10 mL) and dried over MgSO₄. The product was purified by flash chromatography over silica gel eluted with 30% EtOAc/hexanes to give 0.9 g (71%) of the title compound as an oil.

¹H-NMR (300 MHz, CDCl₃, ppm): δ 0.82-0.91 (m, 6H), 0.96 (d, 3H, J=6.8 Hz), 1.52 (m, 2H), 1.88 (m, 1H), 2.67 (br t, 2H), 3.35 (s, 3H), 3.80 (s, 6H), 3.90 (d, 2H, J=6.6 Hz), 6.87 (d, 4H, J=8.8 Hz), 7.20 (d, 4H, J=8.8 Hz), 7.61 (m, 1H), 7.78 (m, 1H), 8.01 (d, 1H, 2H).

Step E: Preparation of 6-(N-methyl-N-isobutyloxycarbonyl)amino-2-propylquinazolin-4(3H)-one

The product of Step D (0.9 g, 1.7 mmol) was added to 3.0 mL of a 10:1 mixture of trifluoroacetic acid and anisole. The solution was stirred for 4 hours, concentrated *in vacuo* and the residue was purified by flash chromatography over silica gel eluted with 50% EtOAc/hexanes to give 0.47 g (88%) of the title compound as a white solid.

¹H-NMR (300 MHz, CDCl₃, ppm): δ 0.89 (d, 6H, J=6.7 Hz), 1.07 (t, 3H, J=7.4 Hz), 1.92 (m, 2H), 2.76 (t, 2H, J=7.8 Hz), 3.40 (s, 3H), 3.93 (d, 2H, J=6.6 Hz), 7.70 (m, 2H), 8.10 (d, 1H, J=2.6 Hz).

Step F: Preparation of methyl 2-(2-chloro-4-hydroxymethyl-6-methoxyphenoxy)-2-phenylacetate

To a solution of 0.500 g (2.65 mmol) of 3-chloro-4-hydroxy-5-methoxybenzyl alcohol (Bader) and 0.668 g (1.1 eq) of methyl 2-

- 80 -

bromophenylacetate dissolved in 5 mL acetone was added 0.733 g (2 eq) of anhydrous potassium carbonate and the reaction mixture was stirred and refluxed overnight. The reaction mixture was cooled to room temperature, filtered and evaporated *in vacuo*. The residual oil was purified on a silica gel flash chromatography column eluted with 35% ethyl acetate/hexane to afford 0.570 g (64%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 1.65-1.75 (t, 1H), 3.70 (s, 3H), 3.80 (s, 3H), 4.55 (d, 2H), 5.75 (s, 1H), 6.80 (s, 1H), 6.90 (s, 1H), 7.30-7.40 (m, 3H), 7.50-7.60 (m, 2H).

FAB-MS: m/e 337, 339 (M+1, 3:1 ratio).

Step G: Preparation of 2-(4-bromomethyl-2-chloro-6-methoxyphenoxy)-2-phenylacetate

To a stirred and cooled (0°C) solution of 0.570 g (1.69 mmol) of the product of Step F dissolved in 6 mL of CH₂Cl₂ was added 0.702 g (2.11 mmol) of carbon tetrabromide and 0.555 g (2.11 mmol) of triphenylphosphine. After the addition the reaction mixture was allowed to warm to room temperature and was stirred 4 hours. The mixture was then evaporated *in vacuo* and purified on a silica gel flash chromatography column eluted with 20% ethyl acetate/hexane to afford 0.580 g (86%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 3.75 (s, 3H), 3.80 (s, 3H), 4.35 (s, 2H), 5.65 (s, 1H), 6.80 (s, 1H), 6.95 (s, 1H), 7.30-7.40 (m, 3H), 7.50-7.60 (m, 2H).

FAB-MS: m/e 398, 400, 402 (M+1).

Step H: Preparation of 3-[4-((1-carbomethoxy-1-phenyl)methoxy)-3-chloro-5-methoxyphenyl]-methyl-6-(N-methyl-N-isobutyloxycarbonyl)-amino-2-propylquinazolin-4(3H)-one

To a half suspension of 80 mg (0.252 mmol) of the product of Step E in 0.5 mL of anhydrous DMF was added 10.6 mg (1.05 eq) of a

- 81 -

60% oil dispersion of sodium hydride and the reaction mixture was stirred under an N₂ atmosphere. After stirring 40 minutes at room temperature, a solution of 0.111 g (1.1 eq) of the product of Step G dissolved in 0.5 mL DMF was added to the solution of the anion. The reaction mixture was then stirred overnight, then partitioned between water and ethyl acetate. The organic layer was separated, dried (MgSO₄), filtered and evaporated *in vacuo*. The residual oil was purified on a silica gel flash chromatography column eluted with 40% ethyl acetate/hexane to afford 100 mg (63%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 0.84-1.01 (m, 9H), 1.58-2.02 (m, 3H), 2.63-2.68 (t, 2H), 3.38 (s, 3H), 3.71 (s, 3H), 3.73 (s, 3H), 3.91 (d, 2H), 5.20-5.30 (br s, 2H), 5.72 (s, 1H), 6.58-6.64 (m, 1H), 6.68 (d, 1H), 7.28-7.34 (m, 3H), 7.48-7.55 (m, 2H), 7.61 (d, 1H), 7.72 (d, 1H), 8.07 (d, 1H).

FAB-MS: m/e 636, 638 (M+1, 3:1 ratio).

Step I: Preparation of 3-[4-((1-carboxy-1-phenyl)-methoxy)-3-chloro-5-methoxyphenyl]methyl-6-(N-methyl-N-isobutyloxycarbonyl)amino-2-propylquinazolin-4(3H)-one

To a solution of 97 mg (0.15 mmol) of the product of Step H dissolved in 2 mL of methanol, was added 0.25 mL of a 1 N solution of NaOH and the reaction mixture was stirred at room temperature for 2.5 hours. The reaction mixture was then adjusted to pH 6 with 1 N HCl, concentrated *in vacuo*, and the residue was applied to a silica gel flash chromatography column and eluted with CHCl₃/MeOH/NH₄OH (80:15:1). Evaporation of the purified fractions and drying *in vacuo* afforded 60 mg (63%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 0.88-1.06 (m, 9H), 1.68-1.82 (m, 2H), 1.85-2.04 (m, 1H), 2.76 (t, 2H), 3.43 (s, 3H), 3.68 (s, 3H), 3.95 (d, 2H), 5.38 (s, 2H), 5.67 (s, 1H), 6.68 (d, 1H), 6.74 (d, 1H), 7.26-7.33 (m, 3H), 7.48-7.54 (m, 2H), 7.72 (d, 1H), 7.82 (dd, 1H), 8.14 (d, 1H).

- 82 -

EXAMPLE 5

3-[4-((1-Carboxy-1-phenyl)methoxy)-3-propylphenyl]methyl-6-(N-methyl-N-iso-butyloxycarbonyl)amino-2-propylquinazolin-4(3H)-one

5 Step A: Preparation of methyl 4-(2-propen-1-yl)-oxybenzoate

A 2 L flask was equipped with a mechanical stirrer, a reflux condenser and a stopper, then charged with 50.05 g (0.329 mol) of methyl 4-hydroxybenzoate, 960 mL of acetone, 22.50 g (1.625 mol) of anhydrous potassium carbonate, 80.5 mL (112.6 g, 0.932 mol) of allyl bromide and the mixture was stirred and refluxed for 14 hours. The mixture was cooled to room temperature, filtered and concentrated to an oil. The residual oil was purified by distillation (97°C @ 0.03 mm Hg) to afford 53.52 g (86%) of the title compound.

15 ¹H NMR (300 MHz, CDCl₃, ppm): δ 3.84 (s, 3H), 4.56 (d, J=7 Hz, 2H), 5.28 (dd, J=3,12 Hz, 1H), 5.40 (dd, J=3,19Hz, 1H), 5.96-6.10 (m, 1H), 6.90 (d, J=10 Hz, 2H), 7.96 (d, J=10 Hz, 2H).

FAB-MS: m/e 193 (M+1).

20 Step B: Preparation of methyl 4-hydroxy-3-(2-propen-1-yl)benzoate

A solution of 15.05 g (78.3 mmol) of the product of Step A in 25 mL of 1,2-dichlorobenzene was magnetically stirred and refluxed (183°C) under an argon atmosphere for 18 hours. At this point, the reaction mixture was cooled to room temperature and applied to a 6 cm diameter by 18 cm silica gel flash chromatography column and eluted with 25% ethyl acetate-hexane to separate the 1,2-dichlorobenzene, then with 40% ethyl acetate-hexane to elute the product. The product fractions were concentrated *in vacuo* and the residual oil was crystallized from hexane to afford 13.70 g (91%) of the title compound.

30 ¹H NMR (300 MHz, CDCl₃, ppm): δ 3.42 (d, J=8 Hz, 2H), 3.88 (s, 3H), 5.14-5.20 (m, 2H), 5.48 (s, 1H), 5.94-6.06 (m, 1H), 6.82 (d, J=12 Hz, 1H), 7.80-7.85 (m, 2H).

- 83 -

FAB-MS: m/e 193 (M+1).

Step C: Preparation of methyl 4-(tert-butyldimethyl-silyloxy)-3-(2-propen-1-yl)benzoate

5 To a solution of 5.168 g (26.9 mmol) of the product of Step B in 50 mL of dichloromethane was added 4.40 mL (2.95 mmol) of triethylamine, 4.46 g (2.95 mmol) of tert-butyldimethylchlorosilane, 0.100 g of 4-dimethylaminopyridine, and the reaction mixture was stirred at room temperature for 2 hours. The mixture was then diluted with 50
10 mL dichloromethane, washed with 100 mL 1 N hydrochloric acid, dried (MgSO₄), filtered and evaporated. The residual oil (7.993 g, 97%) was used in the next step without further purification.

¹H NMR (300 MHz, CDCl₃, ppm): δ 0.24 (s, 6H), 1.02 (s, 9H), 3.36 (d, J=8 Hz, 2H), 3.84 (s, 3H), 4.98-5.08 (m, 2H), 5.88-6.03 (m, 1H), 6.78 (d, J=11 Hz, 1H), 7.76-8.40 (m, 2H).
15

FAB-MS: m/e 307 (M+1).

Step D: Preparation of 4-(tert-butyldimethylsilyloxy)-3-(2-propen-1-yl)benzyl alcohol

20 To a magnetically stirred solution of 8.523 g (28.0 mmol) of the product from Step C in 35 mL of anhydrous THF was added 15.0 mL of a 1.0 M solution of lithium aluminum hydride in THF, and the reaction mixture was stirred under a nitrogen atmosphere for 2 hours. At this
25 point, the reaction was quenched by cautious addition of 10 mL water, the resulting precipitate was dissolved by addition of 1.0 N hydrochloric acid and the product was extracted into ethyl acetate. The organic layer was separated, dried (MgSO₄), filtered and evaporated *in vacuo* to afford 7.258 g (93%) of the title compound.

30 ¹H NMR (300 MHz, CDCl₃, ppm): δ 0.20 (s, 6H), 1.00 (s, 9H), 3.34 (d, J=8 Hz, 2H), 3.84 (s, 1H), 4.57 (s, 2H), 4.97-5.07 (m, 2H), 5.88-6.03 (m, 1H), 6.86 (d, J=10 Hz, 1H), 7.05-7.14 (m, 2H).

- 84 -

FAB-MS: m/e 279, 261 (M+1).

Step E: Preparation of 4-hydroxy-3-(2-propen-1-yl)benzyl alcohol

To a solution of approximately 7.26 g (2.6 mmol) of the crude product of Step D, dissolved in 50 mL of anhydrous THF was added 26 mL (2.6 mmol) of tetra-n-butylammonium fluoride and the reaction mixture was stirred at room temperature for 16 hours. The mixture was then evaporated *in vacuo* and the residual oil was purified on a silica gel flash chromatography column eluted with 5% methanol/chloroform to afford 3.386 g (79%) of the title compound as a colorless oil.

¹H NMR (300 MHz, CDCl₃, ppm): δ 2.12 (br s, 1H), 3.35 (d, J=8 Hz, 2H), 4.54 (s, 3H), 5.05-5.15 (m, 2H), 5.90 (br s, 1H), 5.90-6.05 (m, 1H), 6.70 (d, J=10 Hz, 1H), 7.02-7.10 (m, 2H).

FAB-MS: m/e 165 (M+1).

Step F: Preparation of 4-hydroxy-3-propylbenzyl alcohol

To a solution of 0.370 g (2.25 mmol) of the product of Step E dissolved in 25 mL of absolute ethanol was added 53 mg of a 5% rhodium on carbon catalyst and the mixture was shaken under a 40 psig pressure of hydrogen on a Parr apparatus. After 30 minutes, the reaction mixture was removed, filtered and evaporated *in vacuo*. The residue was purified on a silica gel flash chromatography column eluted with 35% ethyl acetate/hexane to afford the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 0.95 (t, J=8 Hz, 3H), 1.55-1.68 (m, 2H), 2.22 (br s, 1H), 2.57 (t, J=8 Hz, 2H), 4.56 (s, 2H), 5.93 (br s, 1H), 6.66 (d, J=10 Hz, 1H), 7.00 (dd, J=2, 10 Hz, 1H), 7.08 (d, J=2 Hz, 1H).

FAB-MS: m/e 167 (M+1).

Step G: Preparation of methyl (4-hydroxymethyl-2-propylphenoxy)-2-phenylacetate

- 85 -

To a solution of 0.484 g (2.91 mmol) of the product of Step F dissolved in 12 mL of acetone were added 0.667 g (2.91 mmol) of methyl 2-bromophenylacetate, 0.804 g (5.82 mmol) of anhydrous K_2CO_3 and the mixture was stirred and heated at reflux for 5 hours. The mixture was then cooled, filtered and evaporated *in vacuo*. The residual oil was purified on a silica gel flash chromatography column eluted with 30% ethyl acetate/hexane to afford 0.756 g (83%) of the title compound.

1H NMR (300 MHz, $CDCl_3$, ppm): δ 0.95 (t, $J=8$ Hz, 3H), 1.58 (br s, 1H), 1.60-1.75 (m, 2H), 2.70 (t, $J=8$ Hz, 2H), 3.68 (s, 3H), 4.57 (m, 2H), 5.62 (s, 1H), 6.68 (d, $J=10$ Hz, 1H), 7.07 (dd, $J=2, 10$ Hz, 1H), 7.16 (d, $J=2$ Hz, 1H), 7.32-7.44 (m, 3H), 7.55-7.60 (m, 2H).

FAB-MS: m/e 315 ($M+1$).

Step H: Preparation of methyl (4-bromomethyl-2-propylphenoxy)-2-phenylacetate

To a stirred (0°C) solution of 0.750 g (2.31 mmol) of the product of Step G, and 0.949 g (2.86 mmol) of carbon tetrabromide dissolved in 7 mL of methylene chloride was added 0.751 g of triphenylphosphine (2.86 mmol) in portions. After the addition was complete, the reaction mixture was stirred and allowed to warm to room temperature over 1 hour. The reaction mixture was then evaporated *in vacuo*, and the residue was purified on a silica gel flash chromatography column eluted with 10% ethyl acetate/hexane to afford 0.703 g (78%) of the title compound.

1H NMR (300 MHz, $CDCl_3$, ppm): δ 0.95 (t, $J=8$ Hz, 3H), 1.60-1.75 (m, 2H), 2.70 (t, $J=8$ Hz, 2H), 3.69 (s, 3H), 4.44 (s, 2H), 5.62 (s, 1H), 6.64 (d, $J=10$ Hz, 1H), 7.12 (dd, $J=2, 10$ Hz, 1H), 7.18 (d, $J=2$ Hz, 1H), 7.34-7.44 (m, 3H), 7.53-7.58 (m, 2H).

Step I: Preparation of 3-[4-((1-carboxmethoxy-1-phenyl)methoxy)-3-propylphenyl]methyl-6-(N-methyl-N-isobutyloxy-carbonyl)amino-2-propylquinazolin-4(3H)-one

- 86 -

To a solution of 115 mg (0.36 mmol) of the product of Step E from Example 4 in 1.5 mL of anhydrous DMF was added 15.0 mg (0.36 mmol) of a 60% oil dispersion of sodium hydride and the reaction mixture was stirred under an N₂ atmosphere. After stirring 45 minutes at room temperature, a solution of 0.137 g (0.36 mmol) of the product of Step H dissolved in 0.5 mL DMF was added to the solution of the anion. The reaction mixture was then stirred an additional hour at room temperature, then partitioned between water and ethyl acetate. The organic layer was separated, dried (MgSO₄), filtered and evaporated *in vacuo*. The residual oil was purified on a silica gel flash chromatography column eluted with 35% ethyl acetate/hexane to afford 0.060 g (49%) of the title compound.

¹H NMR (300 MHz, CDCl₃, ppm): δ 0.85 (d, 6H), 0.92 (t, 3H), 0.96 (t, 3H), 1.55-2.00 (m, 5H), 2.60-2.75 (m, 4H), 3.36 (s, 3H), 3.77 (s, 3H), 3.90 (d, 2H), 5.28 (br s, 2H), 5.57 (s, 1H), 6.63 (d, 1H), 6.85 (dd, 1H), 6.99 (d, 1H), 7.30-7.42 (m, 3H), 7.50-7.60 (m, 2H), 7.55 (dd, 1H), 7.60 (d, 1H), 8.08 (d, 1H).

FAB-MS: m/e 614 (M+H).

Step J: Preparation of 3-[4-((1-carboxy-1-phenyl)methoxy)-3-propylphenyl]methyl-6-(N-methyl-N-isobutyloxycarbonyl)-amino-2-propylquinazolin-4(3H)-one

To a solution of 60 mg (0.098 mmol) of the product of Step I dissolved in 2 mL of methanol, was added 0.25 mL of a 1 N solution of NaOH and the reaction mixture was stirred at room temperature for 4 hours. The reaction mixture was then adjusted to pH 7 with 1 N HCl, concentrated *in vacuo*, and the residue was applied to a silica gel flash chromatography column and eluted with 10% MeOH/CHCl₃. Evaporation of the purified fractions and drying *in vacuo* afforded 33 mg (57%) of the title compound.

¹H NMR (300 MHz, CD₃OD, ppm): δ 0.88-1.05 (complex, 12 H), 1.56-1.67 (m, 2H), 1.72-1.84 (m, 2H), 1.87-2.00 (m, 1H), 2.55-2.65 (m, 1H).

- 87 -

2.75-2.86 (m, 3H), 3.42 (s, 3H), 3.95 (d, 2H), 5.38 (br s, 2H), 5.43 (s, 1H), 6.85 (d, 1H), 6.94 (dd, 1H), 7.05 (d, 1H), 7.28-7.38 (m, 3H), 7.62-7.70 (m, 2H), 7.71 (d, 1H), 7.83 (dd, 1H), 8.15 (d, 1H).

FAB-MS: m/e 600 (M+H).

5

EXAMPLE 6

3-[4-(1-Carboxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropyl-phenyl]methyl(8-methyl-2-iso-propyl)quinazolin-4(3H)-one

10

Step A: Preparation of 8-methyl-2-iso-propylquinazolin-4-one and 2-(iso-butyryl)-amino-3-methylbenzamide

Dimethylaminopyridine (15 mg) followed by triethylamine (4.14 mL, 0.0279 mol) were added to a stirred solution of 2-amino-3-methylbenzoic acid (1.50 g, 0.0099 mol) in DMF (15 mL) under nitrogen at 0 °C. The *iso*-butyryl chloride (2.07 mL, 0.0198 mol) was added dropwise then the mixture stirred at 0 °C for 30 min. followed by 30 min. at room temperature then at 120 °C for 1 hr. The suspension was then stirred at room temperature for 2 days. Ammonium carbonate (3.5 g) was added in portions and the mixture heated at 120 °C for 3 hr. After cooling to room temperature the mixture was poured into ice/water and extracted with ethyl acetate (3 times). The combined organic phase was washed with water, brine, dried (magnesium sulfate) and the solvent removed *in vacuo* to leave a cream solid. Recrystallisation (1:1 ethyl acetate/hexane) afforded 2-(*iso*-butyryl)amino-3-methylbenzamide (739 mg, 34%).

15

20

25

¹H-NMR 400 MHz (CD₃OD): δ 7.38 (t, 2H), 7.25 (t, 1H), 2.68 (quin, 1H), 2.24 (s, 3H), 1.22 (d, 6H);

30 FAB-MS: 221 (M+H).

The filtrate was evaporated *in vacuo* and chromatographed on silica gel (25, 30% ethyl acetate/hexane) to give 8-methyl-2-iso-propylquinazolin-4-one (340 mg, 17%).

- 88 -

¹H-NMR (CD₃OD): δ 7.99 (dd, 1H), 7.62 (m, 1H), 7.32 (t, 1H), 2.91 (quin, 1H), 2.58 (s, 3H), 1.35 (d, 6H);

FAB-MS: 203 (M+1).

5 Step B: Preparation of 8-methyl-2-iso-propylquinazolin-4-one

5N Sodium hydroxide solution (0.56 mL, 0.0028 mol) was added to a mixture of 2-(iso-butyl)amino-3-methylbenzamide (0.62 g, 0.0028 mol) in methanol (6 mL) and water (14 mL) at room temperature. The mixture was heated at reflux for 1 hr. then cooled to room temperature and acidified with 2N hydrochloric acid solution. Saturated sodium dihydrogen orthophosphate solution was added and the mixture extracted with methylene chloride (4 times). The combined organic phase was washed with water, brine, dried (magnesium sulfate) and the solvent removed *in vacuo* to leave 8-methyl-2-iso-propylquinazolin-4-one (0.43 g, 76%) identical with the material isolated in Step A.

15 Step C: Preparation of methyl 2-bromo-2-(3,4-methylenedioxy-phenyl)acetate

20 A mixture of (3,4-methylenedioxyphenyl)acetic acid (4.64 g, 25.74 mMol) in dry DMF (40 mL), cesium carbonate (9.2 g, 25.74 mMol) and methyl iodide (3.7 g, 26.0 mMol) in dry DMF (40 mL) was stirred at room temperature for 3h. At the end of this period, the reaction mixture was poured into ice water and extracted with ethyl acetate. The organic phase was washed with saturated NaHCO₃, water, brine and then dried (MgSO₄) and filtered. The filtrate was concentrated *in vacuo* to provide pure methyl (3,4-methylenedioxyphenyl)acetate as an oil (4.38 g).

25 N-Bromosuccinimide (3.95 g, 22.2 mMol) and AIBN (0.098 g, 0.06 mMol) were added to a solution of methyl (3,4-methylenedioxy-phenyl)acetate (3.9 g, 21.2 mMol) and the mixture was refluxed for 2.5 h. The reaction was cooled and filtered. The filtrate was concentrated *in vacuo* and the residue obtained was purified by flash chromatography on silica-gel using 10% ethyl acetate-hexane. Yield 2.6 g (oil).

- 89 -

¹H-NMR (400 MHz, CDCl₃, ppm): δ 7.105 (d, 1H), 6.93 (d, 1H), 6.72 (m, 1H), 5.964 (s, 2H), 5.28 (s, 1H), 3.76 (s, 3H).

5 Step D: Preparation of 4-(1-carbomethoxy-1-(3,4-methylenedioxy-phenyl)methoxy)-3,5-dipropylbenzylalcohol

To a solution of (3,5-dipropyl-4-hydroxy)benzyl alcohol (0.19 g, 1.0 mMol) in dry DMF (4 mL) were added cesium carbonate (0.33g, 1.01 mMol) and methyl 2-bromo-2-(3,4-methylenedioxy-phenyl)acetate (0.272 g, 1.0 mMol) and the mixture was stirred at room
10 temperature for 3h. At the end of this period, the reaction mixture was poured into ice water and extracted with ethyl acetate. The organic phase was washed with water, brine, dried (MgSO₄) and then filtered. The filtrate was concentrated *in vacuo* to provide an oil, which was then
15 purified by flash chromatography on silica-gel using ethyl acetate-hexane (1:4) to provide the titled product as a thick colorless oil (0.30 g).

¹H-NMR (200 MHz, CDCl₃, ppm): δ 7.05 (s, 1H), 6.97 (s, 2H), 6.88 (d, 1H), 6.75 (d, 1H), 5.97 (s, 2H), 5.00 (s, 1H), 4.55 (s, 2H), 3.74 (s, 3H), 2.38 (m, 4H), 1.45 (m, 4H), 0.82 (t, 6H).

20

Step E: Preparation of 4-(1-carbomethoxy-1-(3,4-methylenedioxy-phenyl)methoxy)-3,5-dipropylbenzylbromide

To a solution of the product of Step D (0.53 g, 1.38 mMol) in dry THF (10 mL) were added Ph₃P (0.49 g, 2.06 mMol), CBr₄ (0.69 g, 2.06 mMol) and CH₃CN (2 mL), and the mixture was stirred at room
25 temperature for 14 h. At the end of this period, the reaction mixture was concentrated *in vacuo* to provide an oil, which was then purified by flash chromatography on silica-gel using ethyl acetate-hexane (1:9) to provide the titled product as a thick colorless oil (0.57 g).

30

¹H-NMR (400 MHz, CDCl₃, ppm): δ 7.04 (d, 1H), 7.00 (s, 2H), 6.87 (dd, 1H), 6.76 (d, 1H), 5.97 (s, 2H), 5.00 (s, 1H), 4.41 (s, 2H), 3.73 (s, 3H), 2.36 (m, 4H), 1.45 (m, 4H), 0.82 (t, 6H).

- 90 -

Step F: Preparation of 3-[4-(1-carbomethoxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropylphenyl]methyl(8-methyl-2-*iso*-propyl)quinazolin-4(3*H*)-one

Cesium carbonate (57 mg, 0.175 mmol) was added to 8-methyl-2-*iso*-propylquinazolin-4-one (17.7 mg, 0.0875 mmol) in DMF (1 mL) at room temperature under nitrogen. After stirring for 15 min., a solution of 4-(1-carbomethoxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropyl-benzylbromide (52.7 mg, 0.114 mmol), from Step E, in DMF (1 mL) was added and the mixture stirred at room temperature for 12 hr. The solvent was removed *in vacuo* and the residue chromatographed on silica gel (8, 15% ethyl acetate/hexane) to give 3-[4-(1-carbomethoxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropylphenyl]methyl(8-methyl-2-*iso*-propyl)quinazolin-4(3*H*)-one (32 mg, 63%).

¹H-NMR 400 MHz (CDCl₃): δ 8.12 (d, 1H), 7.56 (d, 1H), 7.31 (t, 1H), 7.01 (d, 1H), 6.82 (dd, 1H), 6.73 (m, 3H), 5.95 (s, 2H), 5.32 (br. s, 2H), 4.96 (s, 1H), 3.71 (s, 3H), 3.06 (quin, 1H), 2.60 (s, 3H), 2.30 (m, 4H), 1.53 - 1.30 (m, 4H), 1.22 (d, 3H), 1.21 (d, 3H), 0.77 (t, 6H).

Step G: Preparation of 3-[4-(1-carboxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropylphenyl]-methyl(8-methyl-2-*iso*-propyl)-quinazolin-4(3*H*)-one

5N Sodium hydroxide solution (0.1 mL) was added to a stirred mixture of 3-[4-(1-carbomethoxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropylphenyl]methyl(8-methyl-2-*iso*-propyl)quinazolin-4(3*H*)-one (13.2 mg, 0.0226 mmol) in methanol (1 mL). A few drops of methylene chloride were added to allow stirring then the mixture was stirred at room temperature for 2 hr. The solution volume was reduced to ~10% *in vacuo* then 5% citric acid solution was added. The mixture was extracted with ethyl acetate (3 times). The combined organic phase was washed with water, brine, dried (magnesium sulfate) and the solvent removed *in vacuo* to leave 3-[4-(1-carboxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropylphenyl]methyl(8-methyl-2-*iso*-propyl)quinazolin-4(3*H*)-one (14.2 mg, 100%).

- 91 -

¹H-NMR 400 MHz (CD₃OD): δ 8.04 (d, 1H), 7.64 (d, 1H), 7.36 (t, 1H), 6.96 (d, 1H), 6.88 - 6.70 (m, 4H), 5.95 (s, 2H), 5.40 (s, 2H), 4.97 (s, 1H), 3.17 (quin, 1H), 2.60 (s, 3H), 2.33 (t, 4H), 1.50 - 1.24 (m, 4H), 1.23 (d, 3H), 1.22 (d, 3H), 0.77 (t, 6H);

5

FAB-MS 593.7 (M+Na), 571.8 (M+H).

EXAMPLE 7

10 3-[4-(1-Carboxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropyl-phenyl]methyl(8-methyl-2-(pent-3-yl))quinazolin-4(3H)-one

The title compound was synthesized in a similar manner to that outlined for Example 6 except that in Step A *iso*-butyryl chloride was replaced by 2-ethylbutyryl chloride.

15

¹H-NMR 400 MHz (CDCl₃): δ 8.15 (d, 1H), 7.58 (d, 1H), 7.32 (t, 1H), 6.91 (s, 1H), 6.78 (s, 2H), 6.72 (m, 2H), 5.96 (s, 2H), 5.32 (br. s, 1H), 4.99 (s, 1H), 2.69 (m, 1H), 2.56 (s, 3H), 2.28 (t, 4H), 1.76 (m, 2H), 1.59 (m, 2H), 1.40 (m, 4H), 0.78 (m, 6H), 0.68 (m, 6H);

20

FAB-MS 621.4 (M+Na), 599.6 (M+H).

EXAMPLE 8

25 3-[4-(1-Carboxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropyl-phenyl]methyl(2,8-dimethyl)quinazolin-4(3H)-one

The title compound was synthesized in a similar manner to that outlined for Example 6 except that in Step A *iso*-butyryl chloride was replaced by acetyl chloride.

30

¹H-NMR 400 MHz (CDCl₃): δ 8.15 (d, 1H), 7.58 (d, 1H), 7.32 (t, 1H), 6.91 (s, 1H), 6.81 - 6.72 (m, 4H), 5.98 (s, 2H), 5.28 (br. s, 2H), 5.00 (s, 1H), 2.59 (s, 3H), 2.51 (s, 3H), 2.26 (t, 4H), 1.40 (m, 4H), 0.78 (m, 6H);

FAB-MS 543.9 (M+H).

- 92 -

EXAMPLE 9

3-[4-(1-Carboxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropyl-phenyl]methyl(8-methyl)quinazolin-4(3H)-one

5 Step A: Preparation of 8-methylquinazolin-4(3H)-one
 2-Amino-3-methylbenzoic acid (1.00 g, 0.0066 mol) and
 formamide (12 mL) were heated together at 130°C for 20 hr. under
 nitrogen. After cooling to room temperature, the solid was filtered off,
10 washed with diethyl ether and dried *in vacuo* to leave 8-
 methylquinazolin-4(3H)-one (1.19 g).

¹H-NMR 400 MHz (CD₃OD): δ 8.06 (m, 2H), 7.68 (d, 1H), 7.41 (t, 1H),
 2.59 (s, 3H);

15 EI-MS 160 (M⁺).

Step B: Preparation of 3-[4-(1-carboxy-1-(3,4-methylenedioxy-phenyl))methoxy-3,5-dipropylphenyl]methyl(8-methyl)-quinazolin-4(3H)-one
 3-[4-(1-carboxy-1-(3,4-methylenedioxyphenyl))methoxy-
20 3,5-dipropylphenyl]methyl(8-methyl)quinazolin-4(3H)-one was
 synthesized from 8-methylquinazolin-4(3H)-one and 3,5-dipropyl-4-(1-
 methoxy-carbonyl-1-(3,4-methylenedioxyphenyl)methoxy)-
 benzylbromide in a similar manner to that outlined in steps C and D of
25 Example 6.

¹H-NMR 400 MHz (CDCl₃): δ 8.16 (d, 1H), 8.12 (s, 1H), 7.60 (d, 1H),
 7.38 (t, 1H), 6.95 (m, 3H), 6.82 (d, 1H), 6.73 (d, 1H), 5.98 (s, 2H), 5.10
 (s, 2H), 5.00 (s, 1H), 2.59 (s, 3H), 2.28 (t, 4H), 1.42 (m, 4H), 0.79 (m,
30 6H);

 FAB-MS 551.7 (M+Na), 529.7 (M+H).

- 93 -

EXAMPLE 10

3-[4-(1-Carboxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropyl-phenyl]methyl(8-bromo-2-*iso*-propyl)quinazolin-4(3*H*)-one

5 The title compound was synthesized in a similar manner to that outlined for Example 6 except that in Step A 2-amino-3-methylbenzoic acid was replaced by 2-amino-3-bromobenzoic acid.

1H-NMR 400 MHz (CDCl₃): δ 8.25 (d, 1H), 8.00 (d, 1H), 7.28 (m, 1H), 6.90 (s, 1H), 6.75 (m, 4H), 5.98 (s, 2H), 5.32 (br. s, 2H), 5.01 (s, 1H),
10 3.06 (m, 1H), 2.28 (t, 4H), 1.41 (m, 4H), 1.29 (d, 6H), 0.78 (t, 6H);

FAB-MS 635.2 (M+H).

EXAMPLE 11

15 3-[4-(1-Carboxy-1-(3,4-methylenedioxyphenyl))methoxy-3,5-dipropyl-phenyl]methyl(8-methyl-2-phenyl)quinazolin-4(3*H*)-one

The title compound was synthesized in a similar manner to that outlined for Example 6 except that in Step A *iso*-butyryl chloride was
20 replaced by benzoyl chloride.

1H-NMR 400 MHz (CD₃OD): δ 8.25 (d, 1H), 7.70 (d, 1H), 7.52 - 7.32 (m, 6H), 6.95 (s, 1H), 6.85 - 6.76 (m, 2H), 6.45 (s, 2H), 5.98 (s, 2H), 5.28 (br. s, 2H), 4.99 (s, 1H), 2.59 (s, 3H), 2.28 (t, 4H), 1.38 (m, 4H), 0.76 (t,
25 6H);


FAB-MS 605.7 (M+H).

30

- 94 -

WHAT IS CLAIMED IS:

1. A compound of structural formula I



or a pharmaceutically acceptable salt thereof, wherein:

R^1 is:

- 20
- (a) H,
- (b) (C_1-C_6) -alkyl, (C_2-C_6) -alkenyl or (C_2-C_6) -alkynyl
- each of which is unsubstituted or substituted with a
- substituent selected from the group consisting of:
- 25
- i) aryl as defined below,
- ii) (C_3-C_7) -cycloalkyl,
- iii) Cl, Br, I, F,
- iv) OH,
- v) NH_2 ,
- vi) $NH(C_1-C_4)$ -alkyl,
- 30
- vii) $N[(C_1-C_4)$ -alkyl] $_2$,
- viii) $NHSO_2R^2$,
- ix) CF_3 ,
- x) $COOR^2$, or

- 95 -

- xi) SO_2NHR^3 ; and
- (c) aryl, wherein aryl is defined as phenyl or naphthyl unsubstituted or substituted with 1 or 2 substituents selected from the group consisting of:
- 5 i) Br, I, Cl, F,
 ii) $(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 iii) $(\text{C}_1\text{-C}_4)\text{-alkoxy}$,
 iv) NO_2
 v) CF_3
10 vi) $\text{SO}_2\text{NR}^3\text{R}^3$,
 vii) $(\text{C}_1\text{-C}_4)\text{-alkylthio}$,
 viii) hydroxy,
 ix) amino,
 x) $(\text{C}_3\text{-C}_7)\text{-cycloalkyl}$,
15 xi) $(\text{C}_3\text{-C}_{10})\text{-alkenyl}$; and
- (d) heteroaryl, wherein heteroaryl is defined as a 5- or 6-membered heteroaromatic moiety selected from the group consisting of thiophene, furan, thiazole, oxazole, pyridine or pyrimidine, which is
- 20 unsubstituted, mono- or disubstituted with substituents selected from the group consisting of:
- i) Cl, Br, I, or F,
 ii) OH,
 iii) SH,
25 iv) NO_2 ,
 v) $\text{C}_1\text{-C}_4\text{-alkyl}$,
 vi) $(\text{C}_2\text{-C}_4)\text{-alkenyl}$,
 vii) $(\text{C}_2\text{-C}_4)\text{-alkynyl}$,
 viii) $(\text{C}_1\text{-C}_4)\text{-alkoxy}$, or
30 ix) CF_3 ,
- (e) $(\text{C}_1\text{-C}_4)\text{-perfluoroalkyl}$,
(f) $-\text{O}-(\text{C}_1\text{-C}_6)\text{-alkyl}$,
(g) $-\text{S}(\text{O})_n-(\text{C}_1\text{-C}_9)\text{-alkyl}$,
(h) $-\text{CONR}^3\text{R}^3$, or

- 96 -

(i) $-\text{NR}^3\text{CO}-\text{O}-(\text{C}_1-\text{C}_4)\text{-alkyl}$; and

n is: 0 to 2; and

5 J is: (a) $-\text{C}(=\text{M})-$, (b) J and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} or (c) J and L are connected together to form a 6-membered aromatic ring containing one nitrogen atom, substituted with R^{5a} , R^{5b} and R^{6b} ; and

10

K is: (a) $-\text{C}(=\text{M})-$, (b) K and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} , or (c) K and L are connected together to form a 6-membered aromatic ring containing one nitrogen atom, substituted on the carbon atoms with R^{5a} , R^{5b} and R^{6b} ; and

15

one of a or b is a double bond in Formula I provided that when J is $-\text{C}(=\text{M})-$ then b is a double bond and when K is $-\text{C}(=\text{M})-$ then a is a double bond.

20

L is the point of attachment of the 6-membered fused aromatic ring optionally containing one nitrogen atom; and

M is: O, S or NR^{11} ; and

25

R^2 is:

- (a) H, or
- (b) $(\text{C}_1-\text{C}_6)\text{-alkyl}$; and

30

R^3 is:

- (a) R^2 ,
- (b) $\text{CH}_2\text{-aryl}$,
- (c) aryl, or

- 97 -

(d) (C₃-C₇)-cycloalkyl; andR⁴ groups are independently:

- 5 (a) H,
(b) (C₁-C₆)-alkyl or (C₂-C₆)-alkenyl each of which is
unsubstituted or substituted with one of the following
substituents,
10 i) -OH,
ii) -O-(C₁-C₄)-alkyl,
iii) -S(O)_n-(C₁-C₄)-alkyl,
iv) -NR³-(C₁-C₄)-alkyl,
v) -NHR³,
vi) -COOR³,
vii) -CONHR³,
15 ix) -CONR³R¹¹, or
x) (C₃-C₇)-cycloalkyl,
(c) (C₃-C₇)-cycloalkyl,
(d) F, Cl, Br, I,
(e) CF₃,
20 (f) -COOR³,
(g) -CONR³R¹¹,
(h) -NR³R¹¹,
(i) -NR³CONR³R¹¹,
(j) -NR³COOR¹¹,
25 (k) -SO₂NR³R¹¹,
(l) -O-(C₁-C₄)-alkyl,
(m) -S(O)_n-(C₁-C₄)-alkyl, or
(n) -NHSO₂R¹¹; and

30 R^{5a} and R^{5b} are independently:

- (a) H,
(b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl,
(c) Cl, Br, I, F,
(d) CF₃, or

- 98 -

- (e) when R^{5a} and R^{5b} are bonded to adjacent carbon atoms, they can be joined to form a phenyl ring;

R^{6a} and R^{6b} are independently:

- 5 (a) H,
 (b) aryl-(C₁-C₄)-alkyl,
 (c) heteroaryl-(C₁-C₄)-alkyl,
 (d) (C₁-C₆)-alkyl, which is unsubstituted or substituted
 with a substituent selected from the group consisting
 10 of: -CON(R³)₂, -heteroaryl, -S(O)_n-R¹⁵, -tetrazol-5-
 yl, -CONHSO₂R¹⁵, -SO₂NH-heteroaryl,
 -SO₂NHCOR¹⁵, -PO(OR²)₂, -PO(OR³)₂, -SO₂NH-
 CN, -NR²COOR¹⁵, -OH, -NH₂, guanidino, (C₁-C₄)-
 15 alkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylamino, (C₁-
 C₄)-dialkylamino, -COOR³, -CONHR³, -O-COR³, or
 aryl,
 (e) -CO-aryl,
 (f) (C₃-C₇)-cycloalkyl,
 20 (g) Cl, Br, I, F,
 (h) -OR¹¹,
 (i) -SH,
 (j) -S(O)_n-(C₁-C₄)-alkyl,
 (k) -COR³,
 (l) -CO₂H,
 25 (m) -CO₂-(C₁-C₄)-alkyl,
 (n) -SO₃H,
 (o) -NR²R¹⁵,
 (p) -NR²COR¹⁵,
 (q) -NR²COOR¹⁵,
 30 (r) -SO₂NHR³,
 (s) -SO₂NR²R³,
 (t) -NO₂,
 (u) -NHSO₂CF₃,

- 99 -

- 5 (v) $-\text{CONR}^3\text{R}^3$,
 (w) $-(\text{C}_1-\text{C}_4)\text{-perfluoroalkyl}$,
 (x) $-\text{COOR}^2$,
 (y) $-\text{SO}_3\text{H}$,
 (z) $-\text{N}(\text{R}^2)\text{SO}_2\text{R}^{15}$,
 (aa) $-\text{NR}^2\text{CONR}^3\text{R}^{15}$,
 (bb) $-\text{OC}(=\text{O})\text{NR}^{15}\text{R}^3$,
 (cc) $-\text{aryl}$, wherein aryl is phenyl or naphthyl, which is
 10 unsubstituted or substituted as defined in $\text{R}^1(\text{c})$,
 (dd) $-\text{NHSO}_2\text{CF}_3$,
 (ee) $-\text{SO}_2\text{NH-heteroaryl}$, wherein heteroaryl is
 unsubstituted or substituted as defined in $\text{R}^1(\text{d})$,
 (ff) $-\text{SO}_2\text{NHCOR}^{15}$,
 (gg) $-\text{CONHSO}_2\text{R}^{15}$,
 15 (hh) $-\text{PO}(\text{OR}^2)_2$,
 (ii) $-\text{tetrazol-5-yl}$,
 (jj) $-\text{CONH}(\text{tetrazol-5-yl})$, or
 (kk) $-\text{SO}_2\text{NHCN}$; and

20

 R^7 is:

- (a) H,
 (b) $(\text{C}_1-\text{C}_6)\text{-alkyl}$, unsubstituted or substituted with:
 25 (i) $-\text{aryl}$,
 (ii) $-(\text{C}_3-\text{C}_7)\text{-cycloalkyl}$,
 (iii) $-\text{NR}^3\text{R}^{11}$,
 (iv) $-\text{morpholin-4-yl}$,
 (v) $-\text{OH}$,
 (vi) $-\text{CO}_2\text{R}^3$, or
 30 (vii) $-\text{CON}(\text{R}^3)_2$,
 (c) aryl , unsubstituted or substituted with a substituent
 selected from the group consisting of:
 i) $(\text{C}_1-\text{C}_4)\text{-alkyl}$,
 ii) $-\text{O}-(\text{C}_1-\text{C}_4)\text{-alkyl}$,

- 100 -

- iii) $-\text{CONR}^3\text{R}^{11}$,
- iv) F, Cl, Br or I, or
- v) $-\text{COOR}^3$;

5 R^8 is:

- (a) H,
- (b) $(\text{C}_1\text{-C}_4)\text{-alkyl}$, wherein alkyl is unsubstituted or substituted with one or two substituents selected from the group consisting of:
 - 10 i) $-\text{OH}$,
 - ii) $-\text{NR}^3\text{R}^{11}$,
 - iii) $-\text{COOR}^3$,
 - iv) $-\text{CONHR}^3$, or
 - v) $-\text{CONR}^3\text{R}^{11}$;

15

R^9 and R^{10} are independently:

- (a) H,
- (b) $(\text{C}_1\text{-C}_6)\text{-alkyl}$, unsubstituted or substituted with $(\text{C}_3\text{-C}_7)\text{-cycloalkyl}$,
- (c) $(\text{C}_2\text{-C}_6)\text{-alkenyl}$,
- 20 (d) $(\text{C}_2\text{-C}_6)\text{-alkynyl}$,
- (e) Cl, Br, F, I,
- (f) $(\text{C}_1\text{-C}_6)\text{-alkoxy}$,
- (g) when R^9 and R^{10} are on adjacent carbons, they can be joined to form a phenyl ring,
- 25 (h) perfluoro- $(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (i) $(\text{C}_3\text{-C}_7)\text{-cycloalkyl}$, unsubstituted or substituted with $(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (j) aryl,
- 30 (k) $(\text{C}_1\text{-C}_6)\text{-alkyl-S(O)}_n\text{-(CH}_2)_n\text{-}$,
- (l) hydroxy- $(\text{C}_1\text{-C}_6)\text{-alkyl}$ or dihydroxy- $(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (m) $-\text{CF}_3$,
- (n) $-\text{CO}_2\text{R}^3$,
- (o) $-\text{OH}$,

- 101 -

- (p) $-\text{NR}^3\text{R}^{11}$,
 (q) $-\text{[(C}_1\text{-C}_6\text{)-alkyl]NR}^3\text{R}^{11}$,
 (r) $-\text{NO}_2$,
 (s) $-(\text{CH}_2)_n\text{-SO}_2\text{-N(R}^3\text{)}_2$,
 5 (t) $-\text{NR}^3\text{CO-(C}_1\text{-C}_4\text{)-alkyl}$, or
 (u) $-\text{CON(R}^3\text{)}_2$;

R¹¹ is:

- (a) H,
 10 (b) $(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 (c) allyl,
 (d) $(\text{C}_3\text{-C}_6)\text{-cycloalkyl}$,
 (e) $(\text{C}_1\text{-C}_4)\text{-acyl}$,
 15 (f) benzyl, or
 (g) phenyl; and


R¹², R¹³, R^{14a} and R^{14b} are independently:

- (a) H,
 (b) F, Cl, Br, or I,
 20 (c) $-\text{NO}_2$,
 (d) $-\text{NH}_2$,
 (e) $-\text{NH(C}_1\text{-C}_4\text{)-alkyl}$,
 (f) $-\text{N[(C}_1\text{-C}_4\text{)-alkyl]}_2$,
 25 (g) $-\text{SO}_2\text{NHR}^3$,
 (h) $-\text{CF}_3$,
 (i) $(\text{C}_1\text{-C}_4)\text{-alkyl}$,
 (j) $-\text{OR}^3$,
 (k) $-\text{S(O)}_n\text{-(C}_1\text{-C}_4\text{)-alkyl}$,
 (l) $-\text{NHCO-(C}_1\text{-C}_4\text{)-alkyl}$,
 30 (m) $-\text{NHCO-O(C}_1\text{-C}_4\text{)-alkyl}$,
 (n) $-\text{CH}_2\text{O-(C}_1\text{-C}_4\text{)-alkyl}$,
 (o) $-\text{O-(CH}_2\text{)}_m\text{-OR}^3$,
 (p) $-\text{CONR}^3\text{R}^{11}$, or
 (q) $-\text{COOR}^3$ and

- 102 -

m is 2, 3, or 4; and

R¹² and R¹³ on adjacent carbon atoms can be joined together to form a ring structure:



10

A represents:

15

20

- a) $-Y-C(R^4)=C(R^4)-$,
- b) $-Y-C(R^4)=N-$,
- c) $-Y-N=C(R^4)-$,
- d) $-Y-[C(R^8)(R^8)]_s-Y-$,
- e) $-Y-C(R^8)(R^8)-C(R^8)(R^8)-$,
- f) $-C(R^4)=C(R^4)-Y-$,
- g) $-N=C(R^4)-Y-$,
- h) $-C(R^8)(R^8)-C(R^8)(R^8)-Y-$, or
- i) $-C(R^4)=C(R^4)-C(R^4)=C(R^4)-$; and

25

s is 1 or 2; and

Y is $-O-$, $-S(O)_n-$ and NR^3 ; and

X is:

30

- (a) $-O-$,
- (b) $-S(O)_n-$,
- (c) $-NR^3-$,
- (d) $-CH_2O-$,
- (e) $-CH_2S(O)_n-$,

- 103 -

- 5 (f) $-\text{CH}_2\text{NR}^3-$,
 (g) $-\text{OCH}_2-$,
 (h) $-\text{NR}^3\text{CH}_2-$,
 (i) $-\text{S}(\text{O})_n\text{CH}_2-$, or
 (j) single bond; and

R¹⁵ is:

- 10 (a) aryl, or
 (b) (C₁-C₄)-alkyl, is unsubstituted or substituted with:
 i) NH_2 ,
 ii) $\text{NH}[(\text{C}_1-\text{C}_4)\text{-alkyl}]$,
 iii) $\text{N}[(\text{C}_1-\text{C}_4)\text{-alkyl}]_2$,
 iv) CO_2H ,
 v) $\text{CO}_2(\text{C}_1-\text{C}_4)\text{-alkyl}$,
 15 vi) OH ,
 vii) SO_3H , or
 viii) SO_2NH_2 ; and

Z is:

- 20 (a) $-\text{CO}_2\text{H}$,
 (b) $-\text{CO}_2\text{R}^{16}$,
 (c) -tetrazol-5-yl,
 (d) $-\text{CONH}(\text{tetrazol-5-yl})$
 (e) $-\text{CONHSO}_2\text{-aryl}$, wherein aryl is phenyl or naphthyl,
 25 which is unsubstituted or substituted as defined in
 R¹(c),
 (f) $-\text{CONHSO}_2-(\text{C}_1-\text{C}_8)\text{-alkyl}$, wherein the alkyl group is
 unsubstituted or substituted with a substituent selected
 from the group consisting of: $-\text{OH}$, $-\text{SH}$, $-\text{O}(\text{C}_1-\text{C}_4)\text{-}$
 30 alkyl , $-\text{S}(\text{C}_1-\text{C}_4)\text{-alkyl}$, $-\text{CF}_3$, Cl , Br , F , I , $-\text{NO}_2$,
 $-\text{CO}_2\text{H}$, $-\text{CO}_2-(\text{C}_1-\text{C}_4)\text{-alkyl}$, $-\text{NH}_2$, $-\text{NH}[(\text{C}_1-\text{C}_4)\text{-}$
 $\text{alkyl}]$, $-\text{N}[(\text{C}_1-\text{C}_4)\text{-alkyl}]_2$; and
 (g) $-\text{CONHSO}_2-(\text{C}_1-\text{C}_4)\text{-perfluoroalkyl}$,


- 104 -

- (h) -CONHSO₂-heteroaryl, wherein heteroaryl is unsubstituted or substituted as defined in R¹(d),
- (i) -CONHSO₂NR³R³,
- (j) -SO₂NHCO-aryl, wherein aryl is phenyl or naphthyl, which is unsubstituted or substituted as defined in R¹(c),
- (k) -SO₂NHCO-(C₁-C₈)-alkyl, wherein the alkyl group is unsubstituted or substituted with a substituent selected from the group consisting of: -OH, -SH, -O(C₁-C₄)-alkyl, -S-(C₁-C₄)-alkyl, -CF₃, Cl, Br, F, I, -NO₂, -CO₂H, -CO₂-(C₁-C₄)-alkyl, -NH₂, -NH[(C₁-C₄)-alkyl], -N[(C₁-C₄)-alkyl]₂,
- (l) -SO₂NHCO-(C₁-C₄)-perfluoroalkyl,
- (m) -SO₂NHCO-heteroaryl, wherein heteroaryl is unsubstituted or substituted as defined in R¹(d),
- (n) -SO₂NHCONR³R³,
- (o) -PO(OH)₂,
- (p) -PO(OR²)₂, or
- (q) -PO(OH)(OR²); and

R¹⁶ is:

- (a) (C₁-C₄)-alkyl,
- (b) CHR¹⁷-O-COR¹⁸,
- (c) CH₂CH₂-N[(C₁-C₂)-alkyl]₂,
- (d) CH₂CH₂-N[CH₂CH₂]₂O,
- (e) (CH₂CH₂O)_y-O-[(C₁-C₄)-alkyl], wherein y is 1 or 2,
- (f) aryl or CH₂-aryl, where aryl is as defined above or optionally substituted with CO₂(C₁-C₄)-alkyl,


- 105 -



R¹⁷ and R¹⁸ independently are: (C₁-C₆)-alkyl or phenyl.

- 106 -

2. A compound of structural formula II



or a pharmaceutically acceptable salt thereof, wherein:

 R^1 is:

- 20
- (a) H,
 - (b) (C_1-C_6) -alkyl, (C_2-C_6) -alkenyl or (C_2-C_6) -alkynyl
each of which is unsubstituted or substituted with a
substituent selected from the group consisting of:
 - i) (C_3-C_7) -cycloalkyl,
 - ii) CF_3 ,
 - 25
 - iii) (C_1-C_4) -alkylthio,
 - iv) (C_1-C_4) -alkoxy,
 - (c) (C_1-C_4) -perfluoroalkyl,
 - (d) $-CONR^3R^3$, or
 - (e) $-NR^3CO-O-(C_1-C_4)$ -alkyl; and
- 30

n is: 0, 1, or 2; and

- 107 -

J is: (a) $-C(=M)-$, or (b) J and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} ;

5 K is: (a) $-C(=M)-$, or (b) K and L are connected together to form a 6-membered aromatic ring substituted with R^{5a} , R^{5b} , R^{6a} and R^{6b} , provided that one and only one of J and K is $-C(=M)-$; and

10 one of a or b is a double bond in Formula I provided that when J is $-C(=M)-$, then b is a double bond and when K is $-C(=M)-$, then a is a double bond;

15 L is: the point of attachment of the 6-membered fused aromatic ring optionally containing one nitrogen atom; and

M is: O, S or NR^{11} ; and

R^2 is:

- 20 (a) H,
(b) (C_1-C_6) -alkyl; and

R^3 is:

- 25 (a) R^2 ,
(b) $-CH_2$ -aryl, or
(c) aryl; and

R^4 groups are independently:

- 30 (a) H,
(b) (C_1-C_6) -alkyl or (C_2-C_6) -alkenyl each of which is unsubstituted or substituted with one of the following substituents,
i) $-OH$,
ii) $-O-(C_1-C_4)$ -alkyl,

- 108 -

- iii) $-S(O)_n-(C_1-C_4)\text{-alkyl}$,
- iv) $-NR^3-(C_1-C_4)\text{-alkyl}$,
- v) $-NHR^3$,
- vi) $-COOR^3$,
- 5 vii) $-CONHR^3$,
- ix) $-CONR^3R^{11}$, or
- x) $(C_3-C_7)\text{-cycloalkyl}$,
- (c) $(C_3-C_7)\text{-cycloalkyl}$,
- (d) F, Cl, Br, I,
- 10 (e) CF_3 ,
- (f) $-COOR^3$,
- (g) $-CONR^3R^{11}$,
- (h) $-NR^3R^{11}$,
- (i) $-NR^3CONR^3R^{11}$,
- 15 (j) $-NR^3COOR^{11}$,
- (k) $-SO_2NR^3R^{11}$,
- (l) $-O-(C_1-C_4)\text{-alkyl}$,
- (m) $-S(O)_n-(C_1-C_4)\text{-alkyl}$, or
- (n) $-NHSO_2R^{11}$; and

20

R^{5a} and R^{5b} are independently:

- (a) H,
- (b) $(C_1-C_6)\text{-alkyl}$, $(C_2-C_6)\text{-alkenyl}$ or $(C_2-C_6)\text{-alkynyl}$,
- (c) Cl, Br, I, F,
- 25 (d) CF_3 , or
- (e) when R^{5a} and R^{5b} are bonded to adjacent carbon atoms, they can be joined to form a phenyl ring;

30

R^{6a} and R^{6b} are independently:

- (a) H,
- (b) $\text{aryl-(}C_1-C_4\text{)-alkyl}$,
- (c) $\text{heteroaryl-(}C_1-C_4\text{)-alkyl}$,
- (d) $(C_1-C_6)\text{-alkyl}$, is unsubstituted or substituted with a substituent selected from the group consisting of:

- 5
10
15
20
25
30
- CON(R³)₂, -heteroaryl, -S(O)_n-R¹⁵, -tetrazol-5-yl,
 - CONHSO₂R¹⁵, -SO₂NH-heteroaryl,
 - SO₂NHCOR¹⁵, -PO(OR²)₂, -PO(OR³)₂, -SO₂NH-
 - CN, -NR²COOR¹⁵, -OH, -NH₂, guanidino, (C₁-C₄)-
 - alkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylamino, (C₁-
 - C₄)-dialkylamino, -COOR³, -CONHR³, -O-COR³, or
 - aryl,
 - (e) -CO-aryl,
 - (f) (C₃-C₇)-cycloalkyl,
 - (g) Cl, Br, I, F,
 - (h) -OR¹¹,
 - (i) -SH,
 - (j) -S(O)_n-(C₁-C₄)-alkyl,
 - (k) -COR³,
 - (l) -CO₂H,
 - (m) -CO₂-(C₁-C₄)-alkyl,
 - (n) -SO₃H,
 - (o) -NR²R¹⁵,
 - (p) -NR²COR¹⁵,
 - (q) -NR²COOR¹⁵,
 - (r) -SO₂NHR³,
 - (s) -SO₂NR²R³,
 - (t) -NO₂,
 - (u) -NHSO₂CF₃,
 - (v) -CONR³R³,
 - (w) -(C₁-C₄)-perfluoroalkyl,
 - (x) -COOR²,
 - (y) -SO₃H,
 - (z) -N(R²)SO₂R¹⁵,
 - (aa) -NR²CONR³R¹⁵,
 - (bb) -OC(=O)NR¹⁵R³,
 - (cc) -aryl, wherein aryl is phenyl or naphthyl, which is
- unsubstituted or substituted as defined in R¹(c),

- 110 -

- (dd) $-\text{NHSO}_2\text{CF}_3$,
(ee) $-\text{SO}_2\text{NH}$ -heteroaryl, wherein heteroaryl is
unsubstituted or substituted as defined in R^1 (d),
(ff) $-\text{SO}_2\text{NHCOR}^{15}$,
5 (gg) $-\text{CONHSO}_2\text{R}^{15}$,
(hh) $-\text{PO}(\text{OR}^2)_2$,
(ii) -tetrazol-5-yl,
(jj) $-\text{CONH}(\text{tetrazol-5-yl})$, or
(kk) $-\text{SO}_2\text{NHCN}$; and

10

 R^7 is:

- (a) H,
(b) $(\text{C}_1\text{-C}_6)$ -alkyl, unsubstituted or substituted with:
15 (i) -aryl,
(ii) $-(\text{C}_3\text{-C}_7)$ -cycloalkyl,
(iii) $-\text{NR}^3\text{R}^{11}$,
(iv) -morpholin-4-yl,
(v) -OH,
(vi) $-\text{CO}_2\text{R}^3$, or
20 (vii) $-\text{CON}(\text{R}^3)_2$,
(c) aryl, unsubstituted or substituted with a substituent
selected from the group consisting of:
i) $(\text{C}_1\text{-C}_4)$ -alkyl,
ii) $-\text{O}-(\text{C}_1\text{-C}_4)$ -alkyl,
25 iii) $-\text{CONR}^3\text{R}^{11}$,
iv) F, Cl, Br or I, or
v) $-\text{COOR}^3$;

25

 R^9 and R^{10} are independently:

30

- (a) H,
(b) $(\text{C}_1\text{-C}_6)$ -alkyl, unsubstituted or substituted with
 $(\text{C}_3\text{-C}_7)$ -cycloalkyl,
(c) Cl, Br, F, I,
(d) $(\text{C}_1\text{-C}_6)$ -alkoxy, or

- 111 -

(e) hydroxy-(C₁-C₆)-alkyl or dihydroxy-(C₁-C₆)-alkyl;

R12, R13, R14a and R14b are independently:

- (a) H,
- (b) F, Cl, Br, or I,
- (c) -NO₂,
- (d) (C₁-C₄)-alkyl,
- (e) -OR³,
- (f) -S(O)_n-(C₁-C₄)-alkyl,
- (g) -NHCO-(C₁-C₄)-alkyl,
- (h) -NHCO-O(C₁-C₄)-alkyl,
- (i) -O-(CH₂)_m-OR³,
- (j) -CONR³R¹¹, or
- (k) -COOR³ and

15

m is 2, 3, or 4; and

R12 and R13 on adjacent carbon atoms can be joined together to form a ring structure:

20



25

A represents:

30

- a) -O-C(R⁴)=C(R⁴)-,
- b) -O-C(R⁴)=N-,
- c) -O-[C(R⁸)(R⁸)]_s-O-,
- d) -C(R⁴)=C(R⁴)-O-,
- e) -N=C(R⁴)-O-, or
- f) -C(R⁴)=C(R⁴)-C(R⁴)=C(R⁴)-; and

s is 1 or 2; and

- 112 -

X is:

- (a) -O-,
- (b) -S(O)_n-, or
- (c) -NR³-; and

5

R¹⁵ is:

- (a) aryl, or
- (b) (C₁-C₄)-alkyl, is unsubstituted or substituted with:
 - i) NH₂,
 - ii) NH[(C₁-C₄)-alkyl],
 - 10 iii) N[(C₁-C₄)-alkyl]₂,
 - iv) CO₂H,
 - v) CO₂(C₁-C₄)-alkyl,
 - vi) OH,
 - vii) SO₃H, or
 - 15 viii) SO₂NH₂; and

Z is:


- (a) -CO₂H,
- (b) -tetrazol-5-yl,
- (c) -CONH(tetrazol-5-yl),
- (d) -CONHSO₂-phenyl or -CONHSO₂-naphthyl, wherein
phenyl or naphthyl is unsubstituted or substituted as
defined in R¹(c), or
- 25 (e) -CONHSO₂-heteroaryl, wherein heteroaryl is as
defined in R¹(d).

20

30

- 113 -

3. A compound of structural formula III:



or a pharmaceutically acceptable salt thereof, wherein:

20 R^1 is:

- (a) H,
- (b) (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl or (C₂-C₆)-alkynyl
each of which is unsubstituted or substituted with a
substituent selected from the group consisting of:
- 25 i) (C₃-C₇)-cycloalkyl,
- ii) CF₃,
- iii) (C₁-C₄)-alkylthio,
- iv) (C₁-C₄)-alkoxy,
- (c) (C₁-C₄)-perfluoroalkyl,
- 30 (d) -CONR³R³, or
- (e) -NR³CO-O-(C₁-C₄)-alkyl; and

n is: 0, 1, or 2; and

- 114 -

 R^2 is:

- (a) H,
- (b) (C₁-C₆)-alkyl; and

5

 R^3 is:

- (a) R^2 ,
- (b) -CH₂-aryl, or
- (c) aryl; and

10

 R^4 groups are independently:

- (a) H,
- (b) (C₁-C₆)-alkyl or (C₂-C₆)-alkenyl each of which is unsubstituted or substituted with one of the following substituents,
 - 15 i) -OH,
 - ii) -O-(C₁-C₄)-alkyl,
 - iii) -S(O)_n-(C₁-C₄)-alkyl,
 - iv) -NR³-(C₁-C₄)-alkyl,
 - v) -NHR³,
 - 20 vi) -COOR³,
 - vii) -CONHR³,
 - ix) -CONR³R¹¹, or
 - x) (C₃-C₇)-cycloalkyl,
- (c) (C₃-C₇)-cycloalkyl,
- (d) F, Cl, Br, I,
- 25 (e) CF₃,
- (f) -COOR³,
- (g) -CONR³R¹¹,
- (h) -NR³R¹¹,
- 30 (i) -NR³CONR³R¹¹,
- (j) -NR³COOR¹¹,
- (k) -SO₂NR³R¹¹,
- (l) -O-(C₁-C₄)-alkyl,
- (m) -S(O)_n-(C₁-C₄)-alkyl, or

- 115 -

(n) $-\text{NHSO}_2\text{R}^{11}$; and R^{5a} is:

- (a) H,
- (b) $(\text{C}_1\text{-C}_6)\text{-alkyl}$, $(\text{C}_2\text{-C}_6)\text{-alkenyl}$ or $(\text{C}_2\text{-C}_6)\text{-alkynyl}$,
- (c) Cl, Br, I, F,
- (d) CF_3 , or
- (e) when R^{5a} and R^{5b} are bonded to adjacent carbon atoms, they can be joined to form a phenyl ring;

 R^{6a} is:

- (a) H,
- (b) aryl- $(\text{C}_1\text{-C}_4)\text{-alkyl}$,
- (c) heteroaryl- $(\text{C}_1\text{-C}_4)\text{-alkyl}$,
- (d) $(\text{C}_1\text{-C}_6)\text{-alkyl}$, is unsubstituted or substituted with a substituent selected from the group consisting of:
 $-\text{CON}(\text{R}^3)_2$, heteroaryl, $-\text{S}(\text{O})_n\text{-R}^{15}$, tetrazol-5-yl,
 $-\text{CONHSO}_2\text{R}^{15}$, $-\text{SO}_2\text{NH-heteroaryl}$,
 $-\text{SO}_2\text{NHCOR}^{15}$, $-\text{PO}(\text{OR}^2)_2$, $-\text{PO}(\text{OR}^3)_2$, $-\text{SO}_2\text{NH-CN}$, $-\text{NR}^2\text{COOR}^{15}$, $-\text{OH}$, $-\text{NH}_2$, guanidino, $(\text{C}_1\text{-C}_4)\text{-alkoxy}$, $(\text{C}_1\text{-C}_4)\text{-alkylthio}$, $(\text{C}_1\text{-C}_4)\text{-alkylamino}$, $(\text{C}_1\text{-C}_4)\text{-dialkylamino}$, $-\text{COOR}^3$, $-\text{CONHR}^3$, $-\text{O-COR}^3$, or aryl,
- (e) $-\text{CO-aryl}$,
- (f) $(\text{C}_3\text{-C}_7)\text{-cycloalkyl}$,
- (g) Cl, Br, I, F,
- (h) $-\text{OR}^{11}$,
- (i) $-\text{SH}$,
- (j) $-\text{S}(\text{O})_n\text{-(C}_1\text{-C}_4)\text{-alkyl}$,
- (k) $-\text{COR}^3$,
- (l) $-\text{CO}_2\text{H}$,
- (m) $-\text{CO}_2\text{-(C}_1\text{-C}_4)\text{-alkyl}$,
- (n) $-\text{SO}_3\text{H}$,

- 116 -

- (o) $-\text{NR}^2\text{R}^{15}$,
 (p) $-\text{NR}^2\text{COR}^{15}$,
 (q) $-\text{NR}^2\text{COOR}^{15}$,
 (r) $-\text{SO}_2\text{NHR}^3$,
 5 (s) $-\text{SO}_2\text{NR}^2\text{R}^3$,
 (t) $-\text{NO}_2$,
 (u) $-\text{NHSO}_2\text{CF}_3$,
 (v) $-\text{CONR}^3\text{R}^3$,
 (w) $-(\text{C}_1-\text{C}_4)\text{-perfluoroalkyl}$,
 10 (x) $-\text{COOR}^2$,
 (y) $-\text{SO}_3\text{H}$,
 (z) $-\text{N}(\text{R}^2)\text{SO}_2\text{R}^{15}$,
 (aa) $-\text{NR}^2\text{CONR}^3\text{R}^{15}$,
 (bb) $-\text{OC}(=\text{O})\text{NR}^{15}\text{R}^3$,
 15 (cc) -aryl, wherein aryl is phenyl or naphthyl, which is
 unsubstituted or substituted as defined in $\text{R}^1(\text{c})$,
 (dd) $-\text{NHSO}_2\text{CF}_3$,
 (ee) $-\text{SO}_2\text{NH-heteroaryl}$, wherein heteroaryl is
 unsubstituted or substituted as defined in $\text{R}^1(\text{d})$,
 20 (ff) $-\text{SO}_2\text{NHCOR}^{15}$,
 (gg) $-\text{CONHSO}_2\text{R}^{15}$,
 (hh) $-\text{PO}(\text{OR}^2)_2$,
 (ii) -tetrazol-5-yl,
 25 (jj) $-\text{CONH}(\text{tetrazol-5-yl})$, or
 (kk) $-\text{SO}_2\text{NHCN}$; and

R^9 and R^{10} are independently:

- (a) H,
 30 (b) $(\text{C}_1-\text{C}_6)\text{-alkyl}$, unsubstituted or substituted with
 $(\text{C}_3-\text{C}_7)\text{-cycloalkyl}$,
 (c) Cl, Br, F, I,
 (d) $(\text{C}_1-\text{C}_6)\text{-alkoxy}$, or
 (e) hydroxy- $(\text{C}_1-\text{C}_6)\text{-alkyl}$ or dihydroxy- $(\text{C}_1-\text{C}_6)\text{-alkyl}$;

- 117 -


R12, R13, R14a and R14b are independently:

- (a) H,
- (b) F, Cl, Br, or I,
- (c) -NO₂,
- (d) (C₁-C₄)-alkyl,
- (e) -OR³,
- (f) -S(O)_n-(C₁-C₄)-alkyl,
- (g) -NHCO-(C₁-C₄)-alkyl,
- (h) -NHCO-O(C₁-C₄)-alkyl,
- (i) -O-(CH₂)_m-OR³,
- (j) -CONR³R¹¹, or
- (k) -COOR³ and

m is 2, 3, or 4; and

R12 and R13 on adjacent carbon atoms can be joined together to form a ring structure:

20



A represents:

- a) -O-C(R⁴)=C(R⁴)-,
- b) -O-C(R⁴)=N-,
- c) -O-[C(R⁸)(R⁸)]_s-O-,
- d) -C(R⁴)=C(R⁴)-O-,
- e) -N=C(R⁴)-O-, or
- f) -C(R⁴)=C(R⁴)-C(R⁴)=C(R⁴)-; and

s is 1 or 2; and

- 118 -

X is:

- (a) -O-,
- (b) -S(O)_n-, or
- (c) -NR³-; and

5

R¹⁵ is:

- (a) aryl, or
- (b) (C₁-C₄)-alkyl, is unsubstituted or substituted with:
 - i) NH₂,
 - ii) NH[(C₁-C₄)-alkyl],
 - 10 iii) N[(C₁-C₄)-alkyl]₂,
 - iv) CO₂H,
 - v) CO₂(C₁-C₄)-alkyl,
 - vi) OH,
 - vii) SO₃H, or
 - 15 viii) SO₂NH₂; and


Z is:

- (a) -CO₂H,
- (b) -tetrazol-5-yl,
- (c) -CONH(tetrazol-5-yl),
- (d) -CONHSO₂-phenyl or -CONHSO₂-naphthyl, wherein
phenyl or naphthyl is unsubstituted or substituted as
defined in R¹(c), or
- 25 (e) -CONHSO₂-heteroaryl, wherein heteroaryl is as
defined in R¹(d).

30

- 119 -

4. The compound as recited in Claim 1 wherein the structural formula is:



and the substituents are as defined in the table below:

20	<u>R¹</u>	<u>R^{6a}</u>	<u>R⁹</u>	<u>R¹⁰</u>	<u>R¹², R¹³, R^{14a}, R^{14b}</u>	<u>Z</u>
	Bu	6-Me	H	Allyl	H, H, H, H	COOH
	Bu	6-Me	H	Pr	H, H, H, H	COOH
	Bu	6-Me	H	Cl	H, H, H, H	COOH
25	Pr	6-N(Me)COOiBu	OMe	Cl	H, H, H, H	COOH
	Pr	6-N(Me)COOiBu	H	Pr	H, H, H, H	COOH
	Pr	6-Me	Pr	Pr	2-Br, 5-Br, 3-OMe, 4-OMe	COOH
	H	8-Me	Pr	Pr	2-Br, 5-Br, 3-OMe, 4-OMe	COOH
30	Me	8-Me	Me	Me	3-OMe, 5-OMe	COOH

- 120 -

	<u>R¹</u>	<u>R^{6a}</u>	<u>R⁹</u>	<u>R¹⁰</u>	<u>R¹², R¹³, R^{14a}, R^{14b}</u>	<u>Z</u>
	H	8-Me	H	Pr	3-OMe, 5-OMe	COOH
	H	8-Me	Pr	Pr	3-OMe, 5-OMe	COOH
5	H	8-Me	H	Pr	3-OMe, 5-OMe	CONHSO ₂ Ph
	H	8-Me	Pr	Pr	3-OMe, 5-OMe	CONHSO ₂ Me
	H	8-Me	H	Pr	3-OMe, 5-OMe	CONHSO ₂ Ph(4-iPr)
	i-Pr	8-Br	H	Pr	3-OMe, 5-OMe	CONHSO ₂ Ph(4-iPr)
10	H	6-PhCONH	Bu	H	3-OMe	COOH
	Me	8-Me	Pr	H	3-OMe	COOH
	Me	8-Me	Pr	H	3-OMe	CONHSO ₂ Ph(4-iPr)
	Me	8-Me	Pr	Pr	3-OMe	COOH
15	H	8-Me	Cl	Cl	3-OMe	COOH
	Me	8-Me	Br	Br	3-OMe	COOH
	Ph	8-Me	Cl	Cl	3-OMe	COOH
20	H	8-Me	Pr	Pr	4-OMe	COOH
	Me	6-Me	Pr	Pr	4-OMe	COOH
	i-Pr	8-Me	Pr	H	4-OMe	COOH
	Bu	6-Me	Cl	Cl	4-OMe	COOH
25	Me	8-Me	Br	Br	4-OMe	COOH
	H	8-Me	Pr	Pr	2-OMe	COOH
	Me	6-Me	Pr	Pr	2-OMe, 3-OMe	COOH
	i-Pr	8-Me	Pr	H	2-OMe	COOH

30

- 121 -

	<u>R¹</u>	<u>R^{6a}</u>	<u>R⁹</u>	<u>R¹⁰</u>	<u>R¹², R¹³, R^{14a}, R^{14b}</u>	<u>Z</u>
	Bu	6-Me	Cl	Cl	2-OMe	COOH
	Me	8-Me	Br	Br	2-OMe	COOH
5	H	8-Me	H	Pr	3-OMe, 4-OMe	COOH
	H	8-Me	Pr	Pr	3-OMe, 4-OMe	COOH
	H	8-Me	H	Pr	3-OMe, 4-OMe	CONHSO ₂ Ph
	H	8-Me	Pr	Pr	3-OMe, 4-OMe	CONHSO ₂ Me
10	H	8-Me	H	Pr	3-OMe, 4-OMe	CONHSO ₂ Ph(4-iPr)
	i-Pr	8-Br	H	Pr	3-OMe, 4-OMe	CONHSO ₂ Ph(4-iPr)
	H	6-PhCONH	Bu	H	2-OMe, 3-OMe	COOH
	Me	8-Me	Pr	H	2-OMe, 3-OMe	COOH
15	Me	8-Me	Pr	H	2-OMe, 3-OMe	CONHSO ₂ Ph(4-iPr)
	Me	8-Me	Pr	Pr	2-OMe, 3-OMe	COOH
	H	8-Me	Cl	Cl	2-OMe, 3-OMe	COOH
20	Me	8-Me	Br	Br	2-OMe, 3-OMe	COOH
	H	8-Me	H	Pr	3,4-methylenedioxy	COOH
	H	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
	H	8-Me	H	Pr	3,4-methylenedioxy	CONHSO ₂ Ph
25	H	8-Me	Pr	Pr	3,4-methylenedioxy	CONHSO ₂ Me
	i-Pr	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
	i-Pr	8-Br	Pr	Pr	3,4-methylenedioxy	COOH
	Ph	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
30	(2-Et)Pr	8-Me	Pr	Pr	3,4-methylenedioxy	COOH

- 122 -

	<u>R¹</u>	<u>R^{6a}</u>	<u>R⁹</u>	<u>R¹⁰</u>	<u>R¹², R¹³, R^{14a}, R^{14b}</u>	<u>Z</u>
	H	8-Me	H	Pr	3,4-methylenedioxy	CONHSO ₂ Ph(4-iPr)
	i-Pr	8-Br	H	Pr	3,4-methylenedioxy	CONHSO ₂ Ph(4-iPr)
5	H	6-PhCONH	Bu	H	3,4-methylenedioxy	COOH
	Me	8-Me	Pr	H	3,4-methylenedioxy	COOH
	Me	8-Me	Pr	H	3,4-methylenedioxy	CONHSO ₂ Ph(4-iPr)
	Me	8-Me	Pr	Pr	3,4-methylenedioxy	COOH
10	H	8-Me	Cl	Cl	3,4-methylenedioxy	COOH
	Me	8-Me	Br	Br	3,4-methylenedioxy	COOH
	H	8-Me	H	Pr	5-Br; 3,4-methylenedioxy	COOH
	H	8-Me	Pr	Pr	5-Br; 3,4-methylenedioxy	COOH
15	H	8-Me	H	Pr	5-Br; 3,4-methylenedioxy	CONHSO ₂ Ph(4-iPr).

5. A method of treating a condition in a mammal, the treatment of which is effected or facilitated by a decrease in endothelin mediated actions, comprising the administration, in an amount that is effective for antagonizing the effect of endothelin, of a compound of Structural Formula I as recited in Claim 1.

6. The method as recited in Claim 1, wherein the condition is selected from the group consisting of: hypertension, pulmonary hypertension, Raynaud's disease, myocardial infarction, angina pectoris, acute renal failure, cerebral infarction, cerebral vasospasm, arteriosclerosis, vascular restenosis, asthma, endotoxic shock, endotoxin-induced multiple organ failure or disseminated intravascular coagulation, or cyclosporin-induced renal failure or hypertension.

7. The method as recited in Claim 6, wherein the mammal is human.

- 123 -

8. The method as recited in Claim 7 comprising a pharmaceutical composition of therapeutically effective amount of the compound of formula I and a pharmaceutically acceptable carrier.

5 9. The method as recited in Claim 7, wherein the condition is hypertension.

10 10. A method of treating cardiovascular disorders by administering to a person in need of such treatment a therapeutically effective amount of a compound of Formula I as recited in Claim 1.

15

20

25

30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/02834

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :A61K 31/505; C07D 239/86, 471/04

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-online structure search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Chemical Abstracts, Volume 119, No. 1 issued 1993 (Columbus, Ohio USA) Ries et al, "Preparation of carboxy phenyl alkyl heterocycles---", see Abstract No. 8825u. DE 4,123,341, issued 21 January 1993.	1-10
X	Chemical Abstracts, volume 115, No. 23, issued 1991 (Columbus, Ohio USA) Greenlee et al, "Preparation of benzyl quinazolines---", see Abstract No. 256214v, WO 91/12001, issued 22 August 1991.	1-10
A	US, A, 5,177,095 (GREENLEE et al) 05 January 1993, see columns 1-2.	1-10

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	&	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

15 JUNE 1994

Date of mailing of the international search report

JUL 01 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

JOHN M. FORD jd

Facsimile No. (703) 305-3230

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/02834

A. CLASSIFICATION OF SUBJECT MATTER: US CL :

514/080, 224.2, 230.5, 234.2, 234.5, 234.8, 249, 258, 260, 267, 259; 544/286, 285, 291, 250, 249, 287, 284, 289, 293, 294, 279, 051, 052, 105, 116, 117.

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

514/080, 224.2, 230.5, 234.2, 234.5, 234.8, 249, 258, 260, 267, 259; 544/286, 285, 291, 250, 249, 287, 284, 289, 293, 294, 279, 051, 052, 105, 116, 117.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/02834

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

- I. Claims 1-9, drawn to J-L or K-L with no additional nitrogen in Class 544-287.
 - II. Claims 1-9, drawn to J-L or K-L with an additional nitrogen in Class 544-279.
 - III. Claims 6 and 10 does not comply with PCT Article 13, as more than one use is claimed, 37 CFR 1.475(d).
1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
 2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
 3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
 4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.